



Popularization of improved maize (*Zea mays* L.) production technology through frontline demonstrations in semi arid zone IVA of Rajasthan

B. L. Meena,¹ R. P. Meena, R. R. Meena² and Bhim Singh^{3*}

Krishi Vigyan Kendra, Post Box No. 8, Rajsamand, Maharana Pratap University of Agriculture and Technology, Udaipur - 313 324 (Rajasthan), INDIA

¹Present Address: Krishi Vigyan Kendra, Jhalawar -326 001 (Rajasthan), INDIA

²Krishi Vigyan Kendra, Jhalawar -326 001 (Rajasthan), INDIA

³College of Horticulture and Forestry, Jhalrapatan, Jhalawar -326 023 (Rajasthan), INDIA

*Corresponding author. E-mail: bhimsingh1@gmail.com

Received: May 23, 2014; Revised received: August 20, 2014; Accepted: December 05, 2014

Abstract: Front line demonstrations (FLD,s) on maize (*Zea mays* L.) were laid down at 661 farmers' fields to demonstrate production potential and economic benefits of improved production technologies comprising high yielding varieties namely PM-3, PM-5, PEHM-2, Bio- 9681 and Mahyco 3765 in Rajsamand district of Semi Arid Zone IVa of Rajasthan state during *kharif* seasons from 2006 to 2012 in rainfed farming situation. The improved production technologies recorded an additional yield ranging from 2.50 to 15.78 qha⁻¹ with a mean of 7.94 qha⁻¹. The per cent increase yield under improved production technologies ranged from 33.17 to 68.16 (PM-3), 14.09 to 59.82 (PM-5), 46.61 to 66.97 (PEHM-2), 55.83 to 92.82 (Bio 9681) and 80.12 (Mahyco 3765) in respective years. The average extension gap, technology gap and technology index were 9.10 qha⁻¹, 5.94 qha⁻¹ and 20.08 per cent, respectively in different varieties of maize. The improved production technologies gave higher benefit cost ratio ranging from 1.28 to 3.00 with a mean of 2.45 compared to local checks (1.10) being grown by farmers under locality. The productivity of maize per unit area could be increased by adopting feasible scientific and sustainable management practices with a suitable variety. Considering the above facts, frontline demonstrations were carried out in a systematic and scientific manner on farmer's field to show the worth of a new variety and convincing farming community about potentialities of improved production management technologies of maize for further adoption by the farming community.

Keywords: Front Line Demonstrations, Maize, Production technology, *Zea mays*

INTRODUCTION

Maize (*Zea mays* L.) is the most important world's leading cereal crop which can be grown in diverse seasons, ecologies and uses and known as queen of cereal due to unparallel productivity among cereal crops. Globally, it is cultivated on more than 160 m ha area across 166 countries having wider diversity of soil, climate, biodiversity and management practices. In India, maize occupies third position both in area and production followed by rice and wheat (Anonymous, 2011) According to latest data (2010-11), it is being cultivated on 8.6 m ha with 80 per cent area during *kharif* season. The current maize production is 21.7 mt with an average productivity of 24.35 tha⁻¹. The productivity of India is just half than the world productivity. In Rajasthan it is grown on 0.97 m ha area with production 1.84 m ton and productivity of 1,888 kgha⁻¹ (Anonymous, 2012-13). During recent years, Rajsamand district of Rajasthan has emerged as the leading one in maize production in the state. The productivity of maize in the district is very low as compared to average national productivity (24.35 qha⁻¹). Lack of suitable high yielding variety as

well as poor knowledge about production practices are ascribed as main reasons for low productivity of maize in the district. The productivity of maize per unit area could be increased by adopting recommended scientific and sustainable management production practices using suitable high yielding varieties namely PM-3, PM-5, PEHM-2, Bio-9681 and Mahyco 3765 (Sain Das *et al.*, 2007, Dhaka, 2010 and Ranawat *et al.*, 2011). Frontline demonstration is the new concept of field demonstration evolved by the Indian Council of Agriculture Research (ICAR) with main objective to demonstrate newly released crop production and protection technologies and its management practices in the farmers' fields under different agro-climatic regions of the country under different farming situations. While demonstrating the technologies in the farmer's fields, the scientists are required to study the factors contributing higher crop production, field constraints of production and thereby generate production data and feedback information. Taking into account the above considerations, frontline demonstrations (FLD,s) were carried out in a systematic manner on farmer's field to show the worth of a new variety and convincing farmers to adopt improved production

management practices of maize for enhancing productivity of maize *Zea mays* L. .

MATERIALS AND METHODS

The study was conducted in farmers' fields to demonstrate production potential and economic benefits of improved technologies in Rajsamand district of Semi Arid Zone IVa of Rajasthan state during *kharif* seasons from 2006 to 2012 in rainfed farming situation. To popularize the improved maize production practices, constraints in maize production were identified through participatory approach. Preferential ranking technique was utilized to identify the constraints faced by the respondent farmers in maize production. Farmers were also asked to rank the constraints they perceive as limiting production factor for maize cultivation in order of preference. Based on top rank farmers problems identified, front line demonstrations were planned and conducted at the farmer's fields under ICAR and Integrated Scheme of Oilseeds, Pulses, Oilpalm and Maize (ISOPOM). In all, 661 full package frontline demonstrations were conducted to convince them about potentialities of improved varieties of maize *viz.*, PEHM-2, PM-3, PM-5, Mahyco 3765 and Bio- 9681 during *kharif* seasons from 2006 to 2012 under rainfed farming condition, in light to medium soils with low to medium fertility status under maize-wheat cropping systems. Each demonstration was conducted in an area of 0.4 ha and adjacent to the farmer's fields in which the crop was cultivated with farmer's practice/ local variety. The package of practices included were improved varieties, seed treatment, maintenance of optimum plant stand, recommended fertilizers dose, plant protection measures especially grass hopper management. The spacing followed was at 0.60 m x 0.25 m sown between third week of June to first week of July during the five years with the seed rate of 25 kg/ha. All the participating farmers were trained on all aspects of maize production management. To study the impact of front line demonstrations, out of 661 participating farmers, a total of 120 farmers were selected as respondent through proportionate sampling. Production and economic data for FLDs and local practices were collected and analyzed. The Extension gap, technology gap and technology index were calculated using the formula as suggested by Samui *et al.* (2000).

Extension gap (qha^{-1}) = Demonstration yield (qha^{-1}) – Yield of local check (qha^{-1}).

Technology gap (qha^{-1}) = Potential yield (qha^{-1}) – Demonstration yield (qha^{-1}).

Technology index (%) = Potential yield (qha^{-1}) – (Demonstration yield / Potential yield) x 100

Knowledge level of the farmers about improved production practices of maize before frontline demonstration implementation and after implementation was measured and compared by applying paired t-test

at 5 per cent level of significance. Further, the satisfaction level of respondent farmers about extension services provided was also measured based on various dimensions like training of participating farmers, timeliness of services, supply of inputs, solving field problems and advisory services rendered, fairness of scientists, performance of variety demonstrated and over all impact of FLDs. The selected respondents were interviewed personally with the help of a pre-tested and well structured interview schedule. Client Satisfaction Index was calculated as developed by Kumaran and Vijayaragavan (2005). The individual obtained scores were calculated by the formula as: Client Satisfaction Index = The individual obtained score/ Maximum score possible

The data thus collected were tabulated and statistically analyzed to interpret the FLD,s results.

RESULTS AND DISCUSSION

Constraints in maize production: Farmer's maize production problems were documented in this study. Preferential ranking technique was utilized to identify the constraints faced by the respondent farmers in maize production. The ranking given by the different farmers are given in Table 1. A perusal of table indicates that lack of suitable high yielding variety (HYV) (85.00%) was given the top most rank followed by Low technical knowledge (81.67%), Grass hopper infestation (73.33%), vagaries of weather (70.83%). Based on the ranks given by the respondent farmers for the different constraints revealed that lack of suitable HYV, low technical knowledge, grass hopper infestation are the major constraints to maize production and followed by wild animals. Other constraints such low or erratic rainfall, stem borer infestation, stem rot, weed infestation, water lodging, marketing and post harvest management were found to reduce maize production. Among all the constraints, low soil fertility got least concerns. Other studies (Hassan *et al.*, 1998; Ouma *et al.*, 2002; Joshi *et al.*, 2005; Dhaka *et al.*, 2010; Ranawat *et al.* 2011; Dhruw *et al.*, 2012; Sreelakshmi *et al.*, 2012) have reported similar problems in maize production.

Performance of FLD: A comparison of productivity levels between demonstrated varieties and local checks is shown in table 4. During the period under study, it was observed that the productivity of maize in Rajsamand district under improved production technologies ranged between 16.78 to 32.78 qha^{-1} with a mean yield of 23.29 qha^{-1} . The productivity under improved technologies varied from 16.90 to 24.57, 19.40 to 27.17, 16.78 to 20.89, 28.75 to 32.78 and 30.62 qha^{-1} for the varieties PM-3, PM-5, PEHM-2, Bio 9681 and Mahyco 3765, respectively as against the yield range between 10.05 to 18.45 with a mean of 15.35 qha^{-1} under farmers local practices and varieties during study period. The additional yield of different varieties under improved production technologies over local practices ranged from 2.60 to

Table 1. Ranks given by farmers for different constraints (n=120).

S. No.	Constraints	Percentage	Ranks
1.	Lack of suitable HYV	85.00	I
2.	Stem rot disease	38.33	VII
3.	Stem borer infestation	48.33	VI
4.	Low soil fertility	30.00	X
5.	Low technical knowledge	81.67	II
6.	Wild animals	54.17	V
7.	Vagaries of weather (Delay onset of monsoon, Early withdrawn of monsoon and mid season drought)	70.83	IV
8.	Weed infestation	33.33	VIII
9.	Grass hopper infestation	73.33	III
10.	Water lodging	31.67	IX
11.	Marketing	29.17	XI
12.	Post harvest management	28.33	XII

Table 2. Comparison between knowledge levels of the respondent farmers about improved farming practices of maize (n=120).

Mean score			Calculated 't' value
Before FLD implementation	After FLD implementation	Mean difference	
37.50	62.50	25.00	8.78*

* Significant at 5% probability level

Table 3. Extent of farmers satisfaction of extension services rendered (n=120).

Satisfaction level	Number	Per cent
Low	28	23.33
Medium	55	45.83
High	37	30.83

15.78 qha⁻¹ with a mean of 7.94 qha⁻¹ in comparison to local practice and varieties. The per cent increase yield under improved production technologies ranged from 33.17 to 68.16 (PM-3), 14.09 to 59.82 (PM-5), 46.61 to 66.97 (PEHM-2), (55.83 to 92.82 (Bio 9681) and 80.12 (Mahyco 3765) in respective years. This increased grain yield with improved production technologies was mainly because of high potential yielding varieties.

The variation in the productivity was also caused unusual delay in sowing in some of the farmer's fields. In fields where delayed sowing was done because of prolonged dry spell in the month of July and delay onset of monsoon rains, the crop growth was restricted. The late sowing crop was subjected to relatively less time span available for plant growth and development. Similar yield enhancement in different crops in front line demonstration has amply been documented by Haque (2000), Tiwari and Saxena (2001), Tiwari *et al.* (2003), Nazrul Islam *et al.* (2004), Hiremath *et al.* (2007), Mishra *et al.* (2009), Tomar *et al.* (2009),

Dhaka *et al.* (2010), Kumar *et al.* (2010) and Sreelakshmi *et al.* (2012). From these results it is evident that performance of improved varieties was found better than the local check under local conditions. Farmers were motivated by results of agro technologies applied in the FLDs trials and it is expected that they would adopt these technologies in the coming years also.

Yield of the front demonstration trials and potential yield of the different varieties of crop was compared to estimate the yield gaps which were further categorized into technology index. The technology gap shows the gap in the demonstration yield over potential yield and it was 5.94 qha⁻¹. The observed technology gap may be attributed to dissimilarities in soil fertility, salinity and erratic rainfall and other vagaries of weather conditions in the area. Hence, to narrow down the gap between the yields of different varieties, location specific recommendation appears to be necessary. Technology index shows the feasibility of the variety at the farmer's field. The lower the value of technology index more is the feasibility.

Table 4 revealed that the technology index value was 20.00. The finding of the present study are in line with the findings of Sawardekar *et al.* (2003), Hiremath and Nagaraju (2009) and Dhaka *et al.* (2010).

The economic feasibility of improved technologies over traditional farmer's practices was calculated depending on the prevailing prices of inputs and output

Table 4. Yield of maize as influenced by improved production technologies and high yielding varieties over local practices in farmer's fields (2006 to 2012).

Year	Variety	Area (ha)	Demo. Nos.	Yield (qha ⁻¹)			Improved technology	Add. yield over local check (qha ⁻¹)	increase in yield over local check (%)	Ext. gap (qha ⁻¹)	Tech. gap (qha ⁻¹)	Tech. Index (%)
				Max.	Min.	Av.						
2006	PM-3	16	40	24.38	13.13	16.90	10.05	6.85	68.16	6.85	8.10	32.40
	PEHM-2	34	85	22.50	13.13	16.78	10.05	6.73	66.97	6.73	8.22	32.88
2007	PM-3	10	25	24.00	10.00	18.52	13.30	5.22	39.25	5.22	6.48	25.92
	PM-5	20	50	24.65	12.20	19.44	13.30	6.14	46.17	6.14	5.56	22.24
2008	PM-5	42	55	35.20	24.10	27.17	17.00	10.17	59.82	10.17	7.83	22.37
	PEHM-2	8	20	35.88	22.00	20.89	17.00	3.89	22.88	3.89	9.11	30.37
2009	Mahyco 3765	13	33	39.24	21.00	30.62	17.00	13.62	80.12	13.62	4.38	12.51
	Bio 9681	16	40	41.14	23.68	32.78	17.00	15.78	92.82	15.78	7.22	18.05
	PM-5	60	134	26.40	13.30	23.10	14.75	8.35	56.61	8.35	1.90	7.60
	PM-3	16	40	27.10	13.90	22.25	14.75	7.50	50.85	7.50	2.75	11.00
2010	PM-3	14	35	30.75	22.50	24.57	18.45	6.12	33.17	6.12	0.43	1.72
	PEHM-2	6	15	31.50	23.00	21.05	18.45	2.60	14.09	2.60	3.95	15.80
2011	Bio 9681	12	30	33.10	23.80	28.75	18.45	10.30	55.83	10.30	11.25	28.13
	PM-3	10	25	29.2	21.1	23.4	18.23	6.1	33.19	6.29	0.56	1.73
2012	PEHM-2	5	12	30.2	22.2	20.8	18.23	2.5	15.11	2.39	3.88	15.2
	Bio 9681	5	12	32.32	21.9	28.3	18.23	10.2	54.88	11.2	10.9	29.23
Average	PM-3	10	25	28.3	19.65	23.68	16.8	8.12	32.65	7.4	0.57	1.8
	Bio 9681	10	25	32.8	22.1	28.4	16.8	11.13	55.2	10.1	11.2	28.2
				30.45	18.13	23.29	15.35	7.94	52.83	9.10	5.94	20.08

Table 5. Cost of cultivation (Rs.ha⁻¹), net returns (Rs.ha⁻¹) and Benefit: Cost ratio of maize as affected by improved production technologies over local practices.

Year	Variety	Total cost of cultivation (Rs.ha ⁻¹)		Gross return (Rs.ha ⁻¹)		Net return (Rs.ha ⁻¹)		Benefit : Cost ratio		Add. cost of cultivation	Add. net returns (Rs.ha ⁻¹)
		Improved technology	Local check	Improved technology	Local check	Improved technology	Local check	Improved technology	Local check		
2006	PM-3	8200	6500	19990	13100	11210	7515	1.56	1.01	1700	3695
	PEHM-2	7315	6910	22900	13800	16362	6960	2.36	0.98	405	9402
2007	PM-3	8873	6995	20210	13608	11337	6613	1.28	0.94	1877	4724
	PM-5	8873	6995	22154	13608	13281	6613	1.50	0.94	1877	6668
2008	PM-5	7413	6873	22007	13770	14594	6897	1.96	0.92	540	7697
	PEHM-2	7238	6873	23400	13770	16162	6897	2.23	1.05	365	9265
2009	Mahyco 3765	7638	6873	24802	13770	17164	6897	2.24	1.12	765	10267
	Bio 9681	7913	6873	26551	13770	18638	6897	2.35	1.20	1040	11741
2010	PM-5	6821	6760	24717	15782	17896	6760	2.62	0.96	61	11136
	PM-3	7073	6960	24289	15782	17216	8822	2.43	0.98	113	8394
2011	PM-3	7185	7120	27615	19926	20430	12806	2.84	1.05	65	7624
	PEHM-2	7284	7120	29214	19926	21930	12806	3.00	1.22	164	9124
2012	Bio 9681	8610	7120	32162	19926	23552	12806	2.73	1.30	1490	10746
	PM-3	7185	7120	27615	19926	20430	12806	2.84	1.05	68	8962
Average	PEHM-2	7284	7120	29214	19926	21930	12806	3	1.22	175	8493
	Bio 9681	8610	7120	32162	19926	23552	12806	2.73	1.3	1498	9230
Average	PM-3	7185	7120	27615	19926	20430	12806	2.84	1.09	69	7250
	Bio 9681	8630	7120	32162	19926	23552	12806	2.73	1.4	1496	8563
Average		7714	7004	26399	16886	18733	9753	2.45	1.10	710	8782

costs (Table 5). It was found that cost of production of maize under improved technologies varied from Rs. 7073 to Rs. 8873 ha⁻¹ in case of PM-3, Rs. 6821 to Rs. 8873 ha⁻¹ for PM-5, Rs. 7238 to Rs. 7315 ha⁻¹ for PEHM-2, Rs. 7913 to Rs. 8610 ha⁻¹ for Bio 9681 and Rs. 7638 ha⁻¹ in case of Mahyco 3765 with an average of Rs. 7726 ha⁻¹ with an average of Rs. 6921 ha⁻¹ in local practice. The additional cost incurred in the improved technologies was mainly due to more costs involved in the cost of improved seed only. Front line demonstrations recorded higher mean gross returns (Rs.24616 ha⁻¹) and mean net return (Rs.16906 ha⁻¹) with higher benefit ratio (2.24) under improved technologies of different improved varieties of maize as compared to local checks. These results are in line with the findings of Gurumukhi and Mishra (2003), Sawardekar *et al.* (2003), Sharma (2003), Hiremath *et al.* (2007), Hiremath and Nagaraju (2009) and Sreelakshmi *et al.* (2012). Further, additional cost of Rs.1345 ha⁻¹ in demonstration has yielded additional net returns of Rs. 805 ha⁻¹ with incremental benefit cost ratio 2.24 suggesting its higher profitability and economic viability of the demonstration. Similar results were also reported by Hiremath and Nagaraju (2009) and Dhaka *et al.* (2010) in maize crops. The results from the present study clearly brought out the potential of improved production technologies in enhancing maize production and economic gains in rainfed farming situations conditions of this region of Rajasthan. Hence, maize production technologies have broad scope for increasing the area and productivity at each and every level.

Increase in knowledge: Knowledge level of respondent farmers on various aspects of improved maize production technologies before conducting the frontline demonstration and after implementation was measured and compared by applying paired t-test. It could be seen from the Table 2 that farmers mean knowledge score had increased by 25.00 after implementation of frontline demonstrations. The increase in mean knowledge score of farmers was observed significantly higher. As the computed value of 't-test' (8.78) was statistically significant at 5 % probability level. The results are at par with Narayanaswamy and Eshwarappa (1998) on pulses crops, Singh and Sharma (2004) on mustard crop, Singh *et al.* (2007) on different crops like soyabean, pigeon pea, black gram and Dhaka *et al.* (2010) on maize crop. It means, there was significant increase in knowledge level of the farmers due to frontline demonstration. This shows positive impact of frontline demonstration on knowledge of the farmers that have resulted in higher adoption of improved farm practices. The results so arrived might be due to the concentrated educational efforts made by the scientists.

Farmer's satisfaction: The extent of satisfaction level of respondent farmers over extension services and performance of demonstrated variety was measured by Client Satisfaction Index (CSI) and results presented in

table 3. It is observed that majority of the respondent farmers expressed medium (45.83 %) to the high (30.83 %) level of satisfaction for extension services and performance of technology under demonstrations whereas, very few (23.33) percent of respondents expressed lower level of satisfaction. The results are in close conformity with the results of Narayanaswamy and Eshwarappa (1998) on pulses crops, Kumaran and Vijayaragavan (2005) on mustard & gram crops and Dhaka *et al.* (2010) on maize crop. The medium to higher level of satisfaction with respect to services rendered, linkage with farmer's and technologies demonstrated *etc.* indicate stronger conviction, physical and mental involvement in the frontline demonstration which in turn would lead to higher adoption. This shows the relevance of frontline demonstration. It indicates that maize grown with low yield are identified by low knowledge, unfavourable attitude towards high yielding varieties, low risk bearers with negative perception of maize production technology. In other wards it may also due to then socio-economic status, lower holdings and unavailability of inputs and credit facilities and to some extent supply and marketing problems. This is a point of concern for research and extension functionaries to disseminate improved maize production technologies for raising the productivity of maize at all the levels.

Conclusion

On the basis of the result obtained in present study it can be concluded that the yield gap between conventional practices and improved production technologies was perceptibly higher, there is urgent need to make stronger extension services for educating the cultivators in the implementation of improved production technology. However, the yield level under FLD was better than the local varieties and performance of these varieties could be further improved by adopting recommended production technologies. Hence, it can be observed that increased yield was due to adoption of high yielding varieties and conducting front line demonstration of proven technologies. Yield potentials of crop can be increased to greater extent. This will subsequently increase the income as well as the livelihood of the farming community. From the above research findings it can be also concluded that the maximum number of the respondents had medium level of knowledge and extent of adoption regarding recommended maize production technology. The study reported lack of suitable HYV as major constraint by the beneficiaries and is ranked first followed by low technical knowledge.

REFERENCES

- Anonymous (2011). Vision 2030. Directorate of Maize Research, Pusa, New Delhi (India).
- Anonymous (2012-13). Rajasthan Agricultural Statistics at a

- Glance. Directorate of Agriculture, Rajasthan, Jaipur.
- Dhaka, B.L., Meena, B.S. and Suwalka, R.L. (2010). Popularization of improved maize production technology through frontline demonstrations in south-eastern Rajasthan. *Journal of Agricultural Sciences*, 1(1):39-42.
- Dhruw, K.S., Sengar, R.S. and Yadav, K.N. (2012). Level of knowledge and adoption about recommended maize production technology. *Agriculture Update*, 7(3&4):311-315.
- Gurumukhi, D.R. and Mishra, S. (2003). Sorghum front line demonstration - A success story. *Agriculture Extension Review*, 15(4):22-23.
- Haque, M.S. (2000). Impact of compact block demonstration on increase in productivity of rice. *Maharashtra Journal of Extension Education*, 19(1):22-27.
- Hassan, R.M., Onyango, R. and Rutto, J.K. (1998). Relevance of Maize Research in Kenya to Maize Production Problems Perceived by Farmers. In: RM Hassan (Ed.): A GIS Application for Research Planning in Kenya. Oxon: CAB International.
- Hiremath, S.M. and Nagaraju, M.V. (2009). Evaluation of front line demonstration trials on onion in Haveri district of Karnataka. *Karnataka Journal of Agricultural Sciences*, 22 (5):1092-1093.
- Hiremath, S.M., Nagaraju, M.V. and Shashidhar, K.K. (2007). Impact of front line demonstrations on onion productivity in farmer's field. Paper presented In: *National Seminar on Appropriate Extension Strategies for Management of Rural Resources*, University of Agricultural Sciences, Dharwad, December 18-20, p.100.
- Joshi, P.K., Singh, N.P., Singh, N.N., Gerpacio, R.V. and Pingali, P.L. (2005). *Maize in India: Production Systems, Constraints, and Research Priorities*. Mexico, D.F.: CIMMYT.
- Kumar, A., Kumar, R., Yadav, V.P.S. and Kumar, R. (2010). Impact assessment of frontline demonstrations of Bajra in Haryana state. *Indian Research Journal of Extension Education*, 10 (1):105-108.
- Kumaran, M., Vijayaragavan, K. (2005). Farmers' satisfaction of agricultural extension services in an irrigation command area. *Indian Journal of Extension Education*, 41 (3&4):8-12.
- Mishra, D.K., Paliwal, D.K., Tailor, R.S., Deshwal, A.K. (2009). Impact of frontline demonstrations on yield enhancement of potato. *Indian Research Journal of Extension Education*, 9 (3):26-28.
- Narayanaswamy, C., Eshwarappa, G. (1998). Impact of front line demonstrations. *Indian Journal of Extension Education*, 34(1&2):14-15.
- Nazrul Islam, M., Rezual Karim Md and Safiqul Islam Q.M. (2004). Economic performance of BARI Mash 1 (Improved variety of Black gram) with traditional variety at farmers fields of Bangladesh. *Asian Journal of Plant Sciences*, 3:247-250.
- Ouma, J.H., De Groote and Gethi, M. (2002). Focused Participatory Rural Appraisal of farmer's perceptions of maize varieties and production constraints in the Moist Transitional Zone in Eastern Kenya. IRMA Socio-Economic Working Paper No. 02-01. Nairobi, Kenya: CIMMYT and KARI.
- Ranawat, Y., Ram H., Sisodiya, S.S. and Punjabi, N.K. (2011). Adoption of improved maize cultivation practices by trained and untrained farmers of KVK, Udaipur. *Rajasthan Journal of Extension Education*, 19:144-147.
- Sain Dass, Singh, K.P. and Yadav, V.K. (2007). Present status and potential of maize hybrids in enhancing the productivity. National Conference on "Doubling Maize Production" organized by IFFCO Foundation, ICAR, DMR, DAC and IFFCL at New Delhi from May 08-09, 2007. p.13-19.
- Samui, S.K., Maitra, S., Roy, D.K., Mondal, A.K. and Saha, D. (2000). Evaluation of front line demonstration on groundnut (*Arachis hypogea* L.) in Sundarbans. *Journal of Indian Society of Coastal Agricultural Research*, 18 (2):180-183.
- Sawardekar, S.V., Dhane, S.S. and Jadhav, B.B. (2003). Front-line demonstration performance of salt-tolerant rice varieties in coastal saline soils. *IRRN*, 28 (1):73-74.
- Sharma, O.P. (2003). Moth bean yield improvement through Front Line Demonstrations. *Agriculture Extension Review*, 15 (5):11-13.
- Singh, D.K., Gautam, U.S. and Singh, R.K. (2007). Study on Yield Gap and Level of Demonstrated Crop Production Technology in Sagar District. *Indian Research Journal of Extension Education*, 7 (2&3):94-95.
- Singh, N. and Sharma, F.L. (2004). Impact of front line demonstration on gain in knowledge about mustard production technology among farmers. *2nd National Extension Education Congress*, May 22-24, 2004. Society of Extension Education, Agra & MPUAT, Udaipur: 56.
- Sreelakshmi, C.H., Sameer Kumar, C.V. and Shivani, D. (2012). Productivity enhancement of pigeon pea (*Cajanus cajan* L.) through improved production technology. *Madras Agricultural Journal*, 99 (4-6):248-250.
- Tiwari, K.B. and Saxena, A. (2001). Economic Analysis of FLD of oil seeds in Chindwara. *Bhartiya Krishi Anusandhan Patrika*, 16 (3&4):185-189.
- Tiwari, R.B., Singh, V. and Parihar, P. (2003). Role of front line demonstration in transfer of gram production technology. *Maharashtra Journal of Extension Education*, 22 (1): 19.
- Tomar, R.K.S., Sahu, B.L., Rupendra, K., Singh and Prajapati, R.K. (2009). Productivity enhancement of black gram (*Vigna mungo* L.) through improved production technologies in farmers' field. *Journal of Food Legumes*, 22(3):202-204.