

Research Article

## Probiotic bacteria as a healthy alternative for fish and biological control agents in aquaculture

### Pankaj Das

Fish Biology Laboratory, Postgraduate Department of Zoology, Darrang College, Tezpur-784001 (Assam), India;

Department of Zoology, Gauhati University, Guwahati-781014 (Assam), India

### Puspakhi Borah

Department of Zoology, Nowgong College (Autonomous), Nagaon, Assam-782001, India

### Ramashankar Bordoloi

Department of Zoology, Gauhati University, Guwahati-781014 (Assam), India

### Amrit Pegu

Department of Zoology, Gauhati University, Guwahati-781014 (Assam), India

### Rupam Dutta

Department of Animal Biotechnology, College of Veterinary Science, AAU, Khanapara, Guwahati-781022 (Assam), India

### Chittaranjan Baruah\*

Fish Biology Laboratory, Postgraduate Department of Zoology, Darrang College, Tezpur-784001 (Assam) India

\*Corresponding author. E-mail: chittaranjan.baruah@gmail.com

### Article Info

<https://doi.org/10.31018/jans.v16i2.5543>

Received: February 20, 2024

Revised: May 03, 2024

Accepted: May 08, 2024

### How to Cite

Das, P. *et al.* (2024). Probiotic bacteria as a healthy alternative for fish and biological control agents in aquaculture. *Journal of Applied and Natural Science*, 16(2), 674 - 689. <https://doi.org/10.31018/jans.v16i2.5543>

### Abstract

Aquaculture is a rapidly expanding sector that contributes significantly to the global food supply, with Asia accounting for 90% of production and India being the second largest aquaculture fish producer. Probiotics are used in aquaculture techniques because of the need for aquatic organism development, higher disease resistance, and feed efficiency. Several potential probiotics have been shown to produce antimicrobial substances that inhibit the growth of harmful microbes and preserve intestinal microecological equilibrium by allowing them to stick to stomach surfaces, thus impairing pathogenic growth. Aquaculture has benefited various industries, pharmaceuticals, and global food security in recent decades. Nevertheless, the stress conditions that aquatic creatures experience in aquaculture cause fish to become less resilient to illness and weaken their immune systems. Therefore, impacts local people's socioeconomic conditions and economic progress in many nations. Several approaches, including conventional practices, artificial chemicals, and antibiotics, have been used to control disease in the aquaculture sector. Hence, alternative techniques are far more important to keep the microbial ecosystem in aquaculture systems healthy. This review presents the current understanding of probiotic use in aquaculture as a healthy alternative for fish and biological control agents in aquaculture and to find out its future direction for research.

**Keywords:** Aquaculture, Fish disease, Growth, Probiotic, Lactic acid bacteria

### INTRODUCTION

Microorganisms linked to advantageous outcomes for the host are referred to as probiotics. Aquaculture has decreased the use of antibiotics due to environmental problems created by these agents such as antibiotic-resistant bacteria strains, which adversely affect other ecosystems (Okeke *et al.*, 2022). The addition of antibiotics to the culture species eliminates pathogenic mi-

croorganisms and helpful microorganisms that are beneficial to the same organism. Consequently, accumulating these chemicals in organisms is not safe for human beings, who are the final consumers. Nowadays, there is a growing trend for people to consume 100% natural foods to live a healthier and longer life. Similarly, environmental care has been driven by countless regulations over time in a variety of environments. Probiotics have been studied for fish disease treatment for several

decades, mostly for commercial purposes (Iorizzo *et al.*, 2022). Microorganisms that benefit the body are replacing the need for growth promoters and antimicrobial drugs. The production of antibacterial compounds by some candidate probiotics inhibits the growth of harmful microorganisms that are harmful to the intestinal microbiome. As a result, they adhere to stomach surfaces, preventing pathogenic growth. To prevent pathogenic growth, fish pathogenic bacteria compete for nutrients and attachment sites in the stomach. When tested for direct antibacterial activity against known pathogens, several probiotics used in aquaculture have shown direct antibacterial activity (Xu *et al.*, 2014; Lin *et al.*, 2020).

Aquaculture is a significant contributor to the worldwide food supply. Asia currently provides approximately 90% of the worldwide aquaculture production, making it one of the sectors with the most rapid expansion. Compared to other sources of animal proteins, fish is more readily available and less expensive in tropical nations (Pegu *et al.*, 2023). The escalating global demand for fish presents both a challenge and an opportunity (Naylor *et al.*, 2021). Several experiments show that beneficial bacteria can enhance aquaculture production in addition to disease-causing bacteria (Allameh *et al.*, 2017). Regarding the significant group of microorganisms used as animal probiotics, lactic acid bacteria (LAB) are the most prevalent. LABs are used in the fish production industry to enhance resistance to pathogens, survivability, protein efficiency ratio, feed conversion efficiency, and digestion. Furthermore, they prevent intestinal disorders and neutralize antinutritional factors present in feeds. They also improve the immune system of fish. When used in aquaculture, LABs can improve bacterial growth and monitoring by improving microbial efficiency (Allameh *et al.*, 2017). Probiotic-fortified meals also assist in promoting fish growth, in addition to increasing the metabolic rate of the fish body.

Today, it is known that probiotics possess antimicrobial properties by affecting the intestinal microbiota, secreting antibacterial substances (bacteriocins and organic acids), competing for essential nutrients that enable pathogen survival, and producing antitoxin effects. They inhibit pathogen adhesion to the intestine, compete with pathogens for essential nutrients, and produce antitoxin effects. Furthermore, probiotics can modulate the immune system, regulate allergic responses, and reduce the proliferation of cancer in mammals. Thus, when provided at specific concentrations and viability, probiotics positively affect host health (Razafindralambo *et al.*, 2023). Probiotics are commonly described as "friendly bacteria," "friendly," or "healthy". Studies have been focused on the characteristics of the microorganisms in the intestinal microbiota

for many years. However, the term "probiotic" has primarily been applied to gram-positive lactic acid bacteria (Hoseinifar *et al.*, 2018), particularly those belonging to *Bifidobacterium*, *Lactobacillus* and *Streptococcus* genera.

As opposed to terrestrial animals, aquatic species' gastrointestinal microbiota is particularly dependent on the external environment due to water flow through their digestive tracts. Therefore, there is a great deal of transient bacterial growth in the intestines due to constant intake of water and food, together with microorganisms that live there. Several bacteria have been identified in the gastrointestinal tracts (GIT) of aquatic animals, including *Salmonella*, *Listeria*, and *Escherichia coli*, which may be pathogenic, but probiotic bacteria and various other microorganisms have also been identified. A number of these bacteria are gram-positive, such as *Bacillus*, *Carnobacterium*, *Enterococcus*, and several *Lactobacillus* species; some of these bacteria are also gram-negative, facultatively anaerobic, such as *Vibrio* and *Pseudomonas*, as well as certain fungi, yeasts, and algae belonging to the genera *Debaryomyces*, *Saccharomyces*, and *Tetraselmis* (El-Saadony *et al.*, 2021). There is an increasing interest in aquaculture probiotics at present, (Loh *et al.*, 2017) proposed extending the definition of these probiotics to include "living microbial additives that benefit the health of hydrobionts and therefore increase productivity." Probiotics are often defined as beneficial microorganisms that may survive in the digestive tract due to their resistance to bile salts and acids. In aquaculture, probiotics are relatively new, but interest in them has grown because they reduce disease. The present review presents the effects of probiotic bacteria on immune system responses, rearing parameters, and inhibitory effects on pathogenic bacteria in fish.

#### **AQUATIC PROBIOTICS- SELECTION AND ADMINISTRATION OF THE RIGHT PROBIOTICS**

It is very important to select the right strains of probiotics because inappropriate strains may lead to undesirable effects on the host if the strain is not right. The primary objective of selecting probiotics is to ensure they are safe and capable of producing desirable benefits. Furthermore, they must maintain their ability during production, manufacturing, distribution, and storage before reaching consumers (Shewale *et al.*, 2014). The characteristics of a good and successful probiotic are outlined below, and they should all have a few things in common.

##### **What makes a good probiotic?**

(Michael, 2014; Torres-Maravilla *et al.*, 2024)

It should be a strain capable of positively affecting the host animal, such as increased growth or disease resistance.

It should be non-pathogenic and non-toxic.

A large number of viable cells should be present, preferably in a large quantity.

It should be capable of surviving and metabolizing in the gut environment, such as being resistant to low pH and organic acids.

It should be stable and viable for longer when stored and used in the field.

As recommended by the United Nations, the following specifications should be considered in selecting and approving a probiotic product (Aly, 2010):

The viability of the probiotics should be demonstrated as efficacious and very stable when they pass through the gastrointestinal tract (GIT).

When the probiotic is present in the host's gut, it should colonize.

Competition between probiotic cultures and pathogenic bacteria for attachment to intestinal surfaces

According to in vitro tests, the probiotic should inhibit pathogenic bacteria.

It is important that probiotics are resistant to other sanitary agents or disinfectants.

The genus and species names should be indicated according to international nomenclature on the label of probiotic products.

Labels should indicate doses and expiration dates.

There should be data indicating that the product is not likely to infect an immunocompromised animal.

## MATERIALS AND METHODS

### METHODS OF ADMINISTRATION OF PROBIOTICS IN AQUACULTURE

Various methods of administering probiotics in aquaculture exist, such as feeding, injection, or direct immersion (Hai et al., 2015), individually or in combination, as presented in the Fig.1.

#### Water additives, feed additives, and injection

A probiotic mixture is most administered by incorporating it into the feed (92.8%), followed by direct incorporation into the water (4.8%), and by incorporating it into live food (1.6%) (Melo et al., 2020). Bacterial or fungal

strains can be mixed with feeding pellets, encapsulated into live feedstock, or administered orally in feed for rearing animals (Das et al., 2017). A variety of probiotics, including bacterial strains, yeasts, and substances extracted from them, are generally provided as feed additives in aquaculture. In a recent study, *Lactobacillus plantarum* CR1T5 was applied as a feed additive to improve black-eared catfish growth performance and immunity (Silarudee et al., 2019). Rainbow trout fry's growth performance and nutrient utilization were enhanced by feed additives such as *Bifidobacterium strains* (Sahandi et al., 2018). Gobi et al. (2016) and Gupta et al. (2016) suggest that probiotics can be added to water as an additive. At 106 CFU/ml, *Vibrio lentus* leads to significant changes in gene expression in sea bass related to immune response, cell proliferation and death, cell adhesion, ROS metabolism, and iron transport (Schaeck et al., 2017). Furthermore, applying the product through injection in addition to the methods mentioned above is also possible. Probiotic injection with *Enterobacter sp.* through intramuscular and intraperitoneal injection of strain C6-6 (LaPatra et al., 2014) enhances the immune system of rainbow trout.

#### Single and combination

Probiotics can be applied singly or in combination in a variety of forms, such as multi-strain probiotics, probiotics that contain plant extract, and probiotics that contain yeast extract. It has been found that the combination of probiotics is more effective than using single probiotics, according to most studies on probiotics. Multi-strain probiotics are highly sensitive to pathogens and active against various aquaculture animals (Pannu et al., 2014). The multi-strain probiotics positively affects the growth and survival of rohu at the hatchling and fry stages, but not at the later stages of larval development (Jha et al., 2014). Compared to the application of *Bacillus megaterium* PTB 1.4 alone or *Pediococcus pentosaceus* E2211 alone as a probiotic to catfish *Clarias sp.*, the combination of *B. megaterium* PTB 1.4 and *Pediococcus pentosaceus* E2211 in the feed showed better results than either of the probiotics alone (Hamka et al., 2020). *Lactobacillus plantarum* N11 and *B. velezensis* H3.1 were both combined with *Nile tilapia* (*Oerochromis niloticus*) to promote a higher survival rate (583.33%) when compared to the single application of *L. plantarum* (54.17%) and *B. velezensis* (41.67%) (Doan et al., 2018). Combined application of *B. coagulans* and *Mentha piperita* in *Catla catla* improved growth performance, nutrient retention, and immunity (Bhatnagar and Saluja, 2019).

#### Dosage of probiotics

Depending on the species, the appropriate probiotic dosage should be administered. Probiotic levels must be determined according to the probiont species, the




Fig. 1. Different routes for administration of probiotic bacteria

fish species, their physiological status, the rearing conditions, and the specific application (Hai, 2015). The growth performance of rainbow trout was improved when probiotic *Bifidobacterium* strains were fed at a level of 107 CFU/g to the trout, but not when 100 CFU/g was fed to the trout (Sahandi et al., 2018). As a result, this suggests that increasing the number of probiotics in feed does not always equate to increased growth. When supplementing functional nutrients into aquafeeds, it is important to consider the dose while adding these nutrients. The weight gain of pabda catfish, *Ompok pabda*, was reduced when more than 0.2% of dietary commercial probiotics was provided to the fish (Chowdhury et al., 2020). A study revealed that a concentration greater than a certain level might result in disturbances in the overall body physiology of the fish as well as disturbances in the metabolic process of carbohydrates and fats.

## EFFECTS OF PROBIOTIC BACTERIA ON THE PERFORMANCE OF FISH

### Effects of probiotics on fish growth

The effects of probiotics on both human health and the environment have been investigated in several studies (Nayak et al., 2010; El-Kady et al., 2022). Aquatic animals are often treated with probiotics such as *Enterobacter*, *Pseudomonas*, *Bifidobacteria*, *Carnobacterium*, *Shewanella*, *Lactobacillus*, *Lactococcus*, *Leuconostoc*, *Bacillus*, *Aeromonas*, *Vibrio*, *Clostridium*, and *Saccharomyces* (Nayak et al., 2010, Hai et al., 2015; Hasan et al., 2020).

It investigated how microbial probiotics affect the growth, nutrient digestibility, enzymes, and intestinal microbiota of Rohu fingerlings (*Labeo rohita*) in their diet (Mohapatra et al., 2012). Fish fed a combination of three probiotics (*Bacillus subtilis*, *Lactococcus lactis*, and *Saccharomyces cerevisiae*) had a higher growth rate, protein efficiency ratio, nutrient digestibility, and feed conversion ratio. Several probiotic-containing diets have been shown to increase the growth rates of tilapia (*Oreochromis niloticus* L.) compared to control diets (Anee et al., 2021). Fish performance was improved by adding probiotics to diets containing yeast. In the intestines of wild rainbow trout, yeast was isolated similarly. When introduced into the digestive tract of cultured rainbow trout, this probiotic significantly accelerated their growth. Compared to yeast and control treatments, no significant growth increase was observed with bacterial mixtures in conjunction with optimum protein levels. The results may be explained by yeast's adaptability to aquatic environments more than bacteria. In addition to probiotic-fortified diets, the best growth response was obtained when the bacterium was used instead of yeast. In a study conducted by Zhang

et al. (2021), gram-positive probiotic bacteria were found to enhance the survival, growth, and speed of marine fish larvae. Two kinds of probiotics on the growth of juvenile dentex (*Dentex dentex*) but found no significant improvement over the control group (Ibrahim et al., 2015).

### Feed conversion ratio (FCR)

Adding probiotics to feed can improve feed utilization, even under stressful conditions (Anee et al., 2021). Nile tilapia (*Oreochromis niloticus* L.) showed the greatest FCR values in a probiotic-containing diet. The feed conversion ratio improved when Spirulina was added to N. tilapia's diet as a probiotic (Reque et al., 2022). Furthermore, *N. tilapia* treated with commercial probiotics showed significantly higher feed conversion efficiency than controls. Maniat et al. (2022) found that shrimp-fed probiotics had a lower feed conversion ratio than controls. In a similar study, Mohapatra et al. (2012) found that probiotic diets reduced FCRs in Rohu fingerlings (*Labeo rohita*).

### Protein efficiency ratio (PER)

Food fortified with probiotics may improve the protein efficiency ratio (PER) or apparent nitrogen utilization (ANU) of animals (Anee et al., 2021). Consequently, protein consumption is optimized, which is the most expensive nutrient in feed. Among other benefits, probiotics improve feed nutritional efficiency under stress conditions (Anee et al., 2021). In Nile tilapia (*Oreochromis niloticus* L.), probiotics significantly improved protein efficiency ratios (Reque et al., 2022).

### Digestibility

The use of probiotics in diets has been found to increase nutrient digestibility (Mohapatra et al., 2012). In addition to improving protein digestibility, probiotics may also increase performance (Anee et al., 2021). The positive effects of probiotics on feed nutrient digestion in fish diets can be attributed to the activation of digestive enzymes in bacteria by probiotics (Zhao et al., 2023). Askarian et al. (2012) state that chitin affects Atlantic salmon's adherent aerobic intestinal microbiota (*Salmo salar* L.). Chitin was also tested for protease, amylase, cellulose, phytase, and chitinase activity. LABs have been reported to produce digestive enzymes such as amylase, lipase, and protease. An increase in nutrient digestibility was observed in Rohu fish (*Labeo rohita*) when different types of probiotics were added to their diets (Mohapatra et al., 2012). In terrestrial animals (such as poultry), using probiotics in a diet has been shown to increase digestibility (Wang et al., 2017). Feeding live yeast to sea bass (*Dicentrarchus labrax*) increased their digestive enzyme activity (Tovar-Ramírez et al., 2010; Perdichizzi et al., 2023). In a study conducted by Wang and Xu, probiotics from *Ba-*

*cillus sp.* were shown to significantly increase digestive enzyme activities in common carp by affecting protease, amylase, and lipase activity (Luo *et al.*, 2020). Probiotic treatment significantly affected amylase and trypsin activity in shrimp after probiotic treatment. Probiotics enhance digestion by increasing the population of beneficial microorganisms and the activity of microbial enzymes, thereby improving feed digestibility and absorption. In addition, they demonstrated that digestion enzyme activity is enhanced by high growth performance.

### Body composition

Probiotics were administered to increase protein and lipid content in Nile tilapia (*Oreochromis niloticus* L.) diets. However, the administration of probiotics had no significant effect on the fish's moisture and ash content. When commercial probiotics were used, no statistically significant differences were observed in carcass moisture, ash, and protein content among the treatments (Noshair *et al.*, 2023). The researchers observed differences in the amount of lipids and gross energy in the carcasses, with the highest value measured in fish fed a control diet. The probiotic diets did not significantly alter the body composition of Nile tilapia (Anee *et al.*, 2021). *Bacillus subtilis* also increased fat content in carcasses but not moisture, ash, or protein. *B. subtilis* also improved fat content, not moisture, ash, or protein. Probiotics did not significantly affect fish body composition or tissue synthesis.

### Immune system response

Nutritional factors and feed additives have been studied since the 1980s about fish's immune system. Several of these additives have been investigated by researchers for their ability to protect fish against stressors or diseases (Kiron *et al.*, 2012). Probiotics have a wide range of benefits, but one of their most important applications is their ability to modulate immune systems (Nayak *et al.*, 2010). Alternative methods have gained popularity in recent years to control disease problems. Antibiotic bacteria, such as LAB, are used to control pathogen populations by enhancing immunity or excluding them competitively. To achieve this goal, aquaculture can enrich larval food, include probiotics in the diet, or add them to the water. Fish's immune systems can be stimulated by probiotics both specifically and non-specifically. Fish can benefit from mono-species or multi-species probiotics by enhancing phagocytic and lysozyme enzyme activity and cytokine production. Moreover, probiotic bacteria increase immunoglobulin cells and acidophilic granulocytes in the fish gut, stimulating the immune system. It is possible to influence the immunomodulatory activity of probiotics by varying the source, the type, the dose, and the duration of supplementation (Nayak *et al.*, 2010).

Studies have shown that fish and shellfish contain several immunostimulants that are beneficial to the body. In aquaculture, autochthonous microbiota may augment immune responses to pathogens living in the water. Microbial cell walls contain muramyl dipeptide, glucans, and lipopolysaccharides. Accordingly, probiotic LAB is considered an immunostimulant in aquatic animals. LAB, as a probiotic stimulates the immune system and improves the quality of the water and nutrition, resulting in a higher rate of larval survival and increased aquaculture production (Yu *et al.*, 2022). *Bacillus* S11 protects shrimp from disease by activating both cellular and humoral immune defenses inside the shrimp's gut and presumably providing competitive exclusion. Dead bacterial cells and yeast-glucan have also been shown to stimulate immune responses in shrimp (*Penaeus monodon*). Yu *et al.* (2022) found that different strains of *Lactobacillus rhamnosus* (JCM 1136) induce immune responses in rainbow trout (*Oncorhynchus mykiss*). Fish-feeding supplements such as probiotics and immunostimulants can prevent infectious diseases. Dietary administration of probiotics can enhance the body's natural immune function. The bacteria may adhere transiently to the gastrointestinal tract and colonize it, increasing antibody production (Dysin *et al.*, 2023). Probiotics have been shown to improve disease resistance and animal growth in animal rearing. Consequently, probiotics affect the immune system responses and bacterial population of aquatic organisms and the environment.

Further, some studies suggest that the fish gut microbiota prevents fish from becoming ill from food poisoning by inhibiting the growth of pathogenic bacteria in the digestive system. These results deserve attention because the digestive tract is a key entry point for pathogenic bacteria (Ringo *et al.*, 2006). A list of different probiotic bacteria with their source/medium/fish tested as well as their role in promoting growth is presented in Table 1.

### USE OF PROBIOTICS IN AQUACULTURE

Using probiotics on aquatic organisms has been promoted by the need for sustainable aquaculture. Initial interest was focused on growth promoters and animal health improvements; however, new areas such as reproduction and stress tolerance, require more research. A variety of applications of probiotics in aquaculture are illustrated in Fig. 2 and Table 2.

#### Growth promoter

Aquaculture uses probiotics to stimulate the growth of cultivated species, but it is unclear whether these products increase appetites or improve digestion. When administered over time, microorganisms colonising gastrointestinal tracts can multiply more rapidly than those expelled. Fish cultured with probiotics exhibit

various benefits when these bacteria adhere to their intestinal mucosa (El-Saadony *et al.*, 2021). Several studies have explored the effect of probiotics on phytoplankton (microalgae), a primary source of aquatic food chains, because it is capable of producing nutrient-producing photosynthetic machinery that higher organisms cannot synthesize on their own, such as polyunsaturated fatty acids and vitamins. One of the best microalgae for aquaculture is *Chaetoceros spp.* a primary source of live food. Their nutritional needs, however, have limited production. In the presence of *Chaetoceros muelleri* microalgae, *Vibrio alginolyticus* C7b probiotic achieves a high density. Since rotifers are small, most cultured aquatic species rely on them for their first live food, since they are more accessible to larvae. Rotifers *Brachionus plicatilis* have grown faster when they have been given brine shrimp nauplii, a lactic acid bacterium commonly fed to rotifers. By adding *Lactococcus casei sp.*, *Pediococcus acidilactici*, and *Lactobacillus lactis sp.* lactis best results were achieved (Contreras & Tapia *et al.*, 2020). Probiotics have been used as growth promoters for edible fish. A probiotic *Streptococcus* strain was fed to Nile tilapia (*Oreochromis niloticus*) for 9 weeks, resulting in increased crude protein and crude lipid content and a weight increase from 0.154 g to 6.164 g. Commercial probiotic products were found to increase weight 115.3% with a 2% concentration (Sîrbu *et al.*, 2022). In addition to swordtails (*Xiphophorus maculatus*, *Xiphophorus helleri*), and guppies (*P. sphenops*, *Poecilia reticulata*), the feed was supplemented with *Streptomyces* and *Bacillus subtilis*, and significant survival and growth were found after 90 and 50 days, respectively.

#### Use of probiotics to prevent various diseases

In the aquaculture industry, antibiotics have been used for a long time to prevent diseases from affecting the crop. In addition, this led to the development of bacteria's resistance mechanisms, and an imbalance in aquatic species' gastrointestinal microbiota, which affected their health (Nakano *et al.*, 2007; Sarker *et al.*, 2023). Currently, the European Union regulates the use of antibiotics in organisms for human consumption. Consumers today prefer natural products, free from additives like antibiotics; disease prevention is more common than disease treatment. The use of probiotics in aquaculture is, therefore a viable alternative for inhibiting pathogens and controlling diseases. The probiotic microorganisms release chemicals that are bactericidal or bacteriostatic against pathogenic bacteria in the intestine of the host, thus constituting a barrier to opportunistic pathogen proliferation. Several factors cause the antibacterial effect, including the production of antibiotics, bacteriocins, siderophores, enzymes (lysozymes, proteases), and/or hydrogen peroxide, as well as the generation of organic acids that alter the

intestinal pH. The viable probiotics administered to tilapia *Oreochromis niloticus*, improved non-specific immune response, as measured by parameters such as lysozyme activity, neutrophil migration, and bactericidal activity, which improved fish resistance to infection by *Edwardsiella tarda* (Taoka *et al.*, 2006; Sîrbu *et al.*, 2022).

An isolated strain of *Carnobacterium sp.* from salmon intestine was administered to rainbow trout and Atlantic salmon, demonstrating *in vitro* antagonism against *Aeromonas hydrophila*, *A. salmonicida*, *Flavobacterium psychrophilum*, *Photobacterium damsela*, and *Vibrio species*. The control of furunculosis in rainbow trout was demonstrated to be effective with dead probiotic cultures containing *Vibrio fluvialis* A3-47S, *A. hydrophila* A3-51 and *Carnobacterium* BA211. It appears that in this particular case, the number of leukocytes was greater than with live bacteria, suggesting that humoral factors were less important than cellular immunity in these preparations of inactivated bacteria (Knobloch *et al.*, 2022). In the case of shrimp, studies have focused on the evaluation of probiotics such as *Bacillus cereus*, *Paenibacillus polymyxa*, and *Pseudomonas sp.* PS-102 biocontrol agents against pathogens of various *Vibrio species*. It has been found that probiotic strains isolated from the gastrointestinal tract of clownfish (*Amphiprion percula*) are effective in inhibiting several pathogens, including *A. hydrophila* and *V. alginolyticus*. The bacteria isolated from clownfish can colonize the intestinal mucus and, therefore can be used as a prophylactic and/or therapeutic agent (Knobloch *et al.*, 2022). Because probiotics generate a density *in vivo* that permits the production of antimicrobial metabolites. Also, it has been shown that concentrations of probiotics containing 106 to 108 cells per cell per unit of dietary fiber promote the growth of healthy microbiota in ornamental fish from the genera *Poecilia* and *Xiphophorus*, decreasing the number of heterotrophic microbes. The use of *V. alginolyticus* strains as probiotics to increase survival and growth of white shrimp (*Litopenaeus vannamei*), as well as probiotics in Ecuadorian shrimp hatcheries, increased production by 35%, whereas antimicrobials decreased production by 94% (Pereira *et al.*, 2022).

#### Improvement in nutrient digestion

Probiotics can improve the digestive processes of aquatic animals. Apart from extracellular enzymes, probiotic strains also produce vitamin, fatty acid, and amino acid growth factors (El-Saadony *et al.*, 2021). Nutrient absorption is increased when probiotics are added to feed (Sîrbu *et al.*, 2022). European bass larvae (*Dicentrarchus labrax*) have been treated with probiotics, including several cases in which they are used as food. A strain of bacteria known as *Debaryomyces hansenii* HF1, which produces spermine and spermidine,

**Table 1.** Showing different probiotic bacteria with their source/medium/fish tested as well as their role in promoting growth

Probiotic	Source organism/Medium/ Fish tested	Growth promoting role	Reference
<i>Lactobacillus acidophilus</i>	Added in rearing water	Enhancing water quality in the culture pond of <i>Clarias gariepinus</i>	Dohail et al. (2009)
<i>B. subtilis</i> , <i>B. licheniformis</i> , and <i>Enterococcus faecium</i>	Supplemented with a traditional meal of Rainbow trout	Enhance fish growth by increasing feed consumption efficiency	Merrifield et al. (2010)
<i>L. sakei</i>	Intestines of <i>Paralichthys olivaceus</i>	Macrophage phagocytic activity " Peroxidase activity	Harikrishnan et al. (2010)
<i>Bacillus</i> NL 110, <i>Vibrio</i> Sp. NE 17	Added in rearing water	Enhancing water quality in the culture pond of <i>Macrobrachium rosenbergii</i>	Mujeeb et al. (2010)
<i>Bacillus cereus</i>	Added to the meal of Juvenile <i>Dentex dentex</i> L	Increases fish growth due to a greater food consumption capacity	Merrifield et al. (2010)
<i>Methylococcus capsulatus</i>	Added to the diet of Atlantic salmon ( <i>Salmo salar</i> )	Enhance growth by supplying abundant protein to <i>Salmo salar</i>	Romarheim et al. (2011)
<i>Bacillus</i> spp.	Intestines of <i>Penaeus monodon</i> and shrimp pond sediment	Increase Growth and survival Beneficial microbial probiotics	Nimrat et al. (2011)
<i>Carnobacterium</i> Sp.	Supplemented to salmon bowel	Promote growth by showing in vitro antagonism against known fish pathogens in rainbow trout and Atlantic salmon	Martinez et al. (2012)
<i>Nitrosomonas Nitrobacter</i>	Applied in culture pond.	Promote growth by Oxidizing ammonia and nitrite in the cultured pond	Kaiser et al. (2013)
<i>Enterococcus faecalis</i>	<i>Oncorhynchus mykiss</i>	Favoring growth, stimulation of the immune system, and protection of diseases	Rodriguez-Estrada et al. (2013)
<i>Bacillus</i> Sp. (HFH4, CPF3) <i>Bacillus licheniformis</i> (CPH6, CPF4)	The gastrointestinal tract of <i>Channa punctatus</i> and <i>Heteropneustes fossilis</i>	Proteolytic, Cellulolytic, and amylolytic enzyme activity	Banerjee et al. (2013)
<i>Lactobacillus mesenteroides</i> SMM69 <i>Weissella cibaria</i> P71	<i>Scophthalmus maximus</i> L.	Antimicrobial activity against the turbot pathogens <i>T. maritimum</i> and <i>V. splendidus</i>	Muñoz-Atienza et al. (2014)
<i>Bacillus subtilis</i> <i>Bacillus licheniformis</i> <i>Bacillus</i> sp. <i>Pediococcus</i> sp.	<i>Oreochromis</i> sp.	Resistance to <i>S. agalactiae</i>	Ng WK et al. (2014)
<i>Lactobacillus mesenteroides</i> SMM69 <i>Weissella cibaria</i> P71	<i>Cophthalmus maximus</i> L.	Antimicrobial activity against the turbot pathogens <i>T. maritimum</i> and <i>V. splendidus</i>	Muñoz-Atienza et al. (2014)
<i>Lactobacillus Plantarum</i> and <i>Pseudomonas</i> fluorescent	Added in pond water	Increase survival rate by 96.22% in <i>Clarias gariepinus</i>	Omenwa et al. (2015)
<i>Bacillus</i> sp. <i>Pediococcus</i> sp.	<i>Solea senegalensis</i>	improvement protection against pathogen outbreaks	Batista et al. (2015)
<i>Lactobacillus plantarum</i> (LP20)	<i>Seriola dumerili</i>	Improves immune response and stress	Dawood et al. (2015)
<i>Lactobacillus rhamnosus</i>	<i>Pagrus major</i>	Growth-promoting agent and Increases growth	Dawood et al. (2016)
<i>Bacillus megaterium</i> PTB 1.4	Catfish	Increased the activity of digestive enzymes and the growth of catfish	Afrilasari et al. (2016)
<i>Bacillus megaterium</i> , <i>Bacillus polymyxa</i> <i>Lactobacillus delbrueckii</i> <i>Lactobacillus acidophilus</i> <i>Bacillus subtilis</i> <i>Lactobacillus bulgaricus</i> <i>Saccharomyces cerevisiae</i>	<i>Oreochromis</i> sp.	Increased the performance of zootechnical parameters	Gutiérrez et al. (2016)
	<i>C. gariepinus</i>	Increases larval survival	Dennis et al. (2016)


Table 1.Cont.

<i>Pseudoalteromonas</i> sp.	Fish	Inhibitory activity against fish pathogens	Sayes et al., 2016
<i>Enterococcus casseliflavus</i>	<i>Oncorhynchus mykiss</i> .	Capability of improving growth performance and enhancing disease resistance by immunomodulation	Safari et al. (2016)
<i>Pseudoalteromonas</i> sp. Ceba MLms gA3	Fish	Inhibitory activity against the pathogen <i>V. anguillarum</i>	Wesseling et al. (2016)
<i>Nitrosomonas</i> Sp. <i>Nitrobacters</i> Sp.	The gut of Catla ( <i>Catla catla</i> ), Rohu ( <i>Labeo rohita</i> ), and Grass carp ( <i>Ctenopharyngodon idella</i> )	Block the pathogen entry in the culture pond and raise the concentration of dissolved oxygen.	Sunitha and Krishna, (2016)
<i>Escherichia coli</i> , <i>Proteus vulgaris</i> , <i>Bacillus subtilis</i> , <i>Klebsiella pneumonia</i> , <i>Pseudomonas aeruginosa</i> and <i>Staphylococcus aureus</i>	<i>Scomberomorus guttatus</i>	-	Karthiga et al. (2016)
<i>Bacillus licheniformis</i> (TSB27) <i>Lactobacillus thuringiensis</i> <i>Bacillus Plantarum</i> <i>Bacillus subtilis</i> (B46). <i>Bacillus subtilis</i> WB60	<i>Sparus aurata</i> L.	Enhances the immune system	Bahi et al. (2017)
<i>Bacillus pumilus</i> H2	<i>Anguilla japonica</i>	Increased in weight, efficiency in food and protein	Lee et al. (2017)
<i>Bacillus pumilus</i> H2	Fish	Anti-Vibrio activity	Gao et al. (2017)
<i>Pseudoalteromonas</i> sp	<i>Seriola lalandi</i>	Increased larval survival	Leyton et al. (2017); Mata et al. (2017)
<i>Lac. pentosus</i> BD6, <i>Lac. fermentum</i> LW2, <i>Bacillus subtilis</i> E20, <i>Saccharomyces cerevisiae</i> P13	Asian seabass	Improved either the growth performance or disease resistance of Asian seabass against <i>A. hydrophila</i>	Lin et al. (2017)
<i>Bacillus subtilis</i> <i>Lactobacillus rhamnosus</i>	<i>Labeo rohita</i>	Increased the value of biochemical components	Munirasu et al. (2017)
<i>Lactobacillus plantarum</i>	<i>Oreochromis niloticus</i>	Decreases mortality and improves growth	Meidong et al. (2017)
<i>Bacillus</i> sp. MVF1	<i>Labeo rohita</i>	Decreased susceptibility to disease	Nandi et al. (2017)
<i>Lactobacillus casei</i>	<i>Tor tambra</i>	Growth performance and feed efficiency increased	Muchlisin et al. (2017)
<i>Kocuria</i> SM1 <i>Rhodococcus</i> SM2	<i>Oncorhynchus mykiss</i>	Produces extracellular enzymes that may have a role in the host digestive processes	Sharifuzzaman et al. (2017)
<i>Bacillus subtilis</i> <i>Bacillus licheniformis</i> <i>Lactobacillus plantarum</i>	Juvenile rainbow trout Tilapia ( <i>Oreochromis</i> sp.)	Resistance against <i>A. salmonicida</i> Enhanced the growth performance and modulated some hematological parameters.	Park et al. (2017) Yamashita et al. (2017)
<i>Vibrio lentus</i>	<i>Dicentrarchus labrax</i>	Protective effect against Vibriosis caused by <i>V. harveyi</i> in sea bass larvae	Schaeck et al. (2017)
<i>Psychrobacter</i> spp	digestive system of Turbot, wild ( <i>Scophthalmus maximus</i> )	antagonistic activity towards pathogen <i>Tenacibaculum maritimum</i>	Wanka et al. (2018)
<i>Clostridium butyricum</i>	Kuruma shrimp ( <i>Marsupenaeus japonicas</i> )	Increased the content of intestinal short-chain fatty acid along with propionic acid and butyric acid and increased the levels of body crude protein.	Duan et al. (2018)
<i>Pseudomonas</i> sp. (GP21) and <i>Psychrobacter</i> sp. (GP12)	The gastrointestinal tract of Atlantic Cod	Elicit the immune response, particularly in mucosal surfaces such as the skin and intestine.	Butt, et al. (2021)
<i>Lactobacillus rhamnosus</i>	Administered to <i>Oreochromis niloticus</i>	Stimulate the phagocytic activity in Tilapia	Cano-Lozano et al. (2022)



**Table 1.Cont.**

<i>Bacillus</i> Sp.	Common carp, <i>Cyprinus carpio</i>	Increase Survival rate Protease activity Lipase activity	Sumon et al. (2022)
<i>Bacillus</i> spp	Added in the surrounding water	Increase growth performance in cultured pond by phosphate solubilization and phosphate production	Armandeh et al. (2022)
<i>Bacillus cereus</i> , <i>Paenibacillus polymyxa</i> , and <i>Pseudomonas</i> Sp. PS-102	-	Act as the antagonist against various pathogens in shrimp	Sanches-Fernandes et al. (2022)
<i>Lactobacillus acidophilus</i> (Fla) and <i>Saccharomyces boulardii</i> (FSB)	Supplemented with the diet of <i>Acipenser baerii</i>	Enhance growth, immunity, and survival rate	Mocanu et al. (2022)
<i>B. coagulans</i> SC8168	Added in rearing water	Enhancing water quality in the culture pond of <i>Penaeus vannamei</i>	Monier et al. (2023)
<i>Bacillus subtilis</i>	Gut of <i>Cirrhinus mrigala</i>	Increase the length and weight of guppies ( <i>Poecilia reticulata</i> , <i>P. sphenops</i> ), and swordtail ( <i>Xiphophorus helleri</i> , <i>X. maculatus</i> ), and increase the proteases and amylases activity in ornamental fish	Hoseinifar et al. (2023)
<i>Debaryomyces hansenii</i> HF1	Supplemented to Sea bass larvae	Promote growth by secreting amylase and trypsin, enzymes that help in digestion	Perdichizzi et al. (2023)
<i>Psychrobacter</i> Sp. (GP11), <i>Shewanella</i> Sp. (GS11), <i>Photobacterium</i> Sp. (GP31), and <i>Vibrio</i> Sp. (GV11),	Atlantic cod's intestine	Differential expression of an immune gene associated with bacterial defense and inflammation in the head kidney leukocytes	Coulibaly et al. (2023)
<i>Bacillus</i> sp.	Added in rearing water	Increasing growth through improving the water quality in the <i>Penaeus monodon</i> culture pond	Torres-Maravilla et al. (2024)



**Fig. 2. Use of probiotics in aquaculture**

grows in the gastrointestinal tracts of mammals. The enzymes secreted by this yeast also aid the digestion of sea bass larvae. 0.5 g of *Bacillus cereus* strain Ekg+1 supplemented in the diet increased the growth of juvenile common dentex *Dentex dentex* L. (Meng et al., 2023). *B. licheniformis*, *B. subtilis*, and *Enterococcus*

*faecium* were also found to produce similar results for rainbow trout when given with a diet that included *Bacillus subtilis*, *B. licheniformis*, and *Enterococcus faecium* for 10 weeks (Merrifield et al., 2010). There have been some studies where European sea bass larvae have been fed a probiotic yeast diet (*Saccharomyces cere-*

*visiae* strain X2180), evaluating growth and antioxidant enzyme activity (glutathione peroxidase, catalase, and superoxide dismutase), indicating that probiotic supplementation may alter enzyme activity and gene expression patterns (Tovar-Ramírez et al., 2010). Various strains of *Bacillus* have been used as probiotics to increase the apparent digestibility of crude protein, dry matter, and phosphorus in white shrimp *Litopenaeus vannamei* Boone and Fen *Neropenaeus indicus*. The organisms' size increased when the diet was supplemented with 50 grams of probiotics per kilogram of food (Lin et al., 2020). To maintain a constant effect on digestive enzyme production, the probiotic should be managed throughout the shrimp's development (Ziaei-Nejad et al., 2006). *Bacillus subtilis*, isolated from the intestine of *Cirrhinus mrigala*, has been incorporated into the diet of guppies (*Poecilia reticulata*, *P. sphenops*), and swordtails (*Xiphophorus helleri*, *X. maculatus*) to assess their intestinal health. A significant increase has been observed in the length and weight of ornamental fish, as well as an increase in the specific activity of proteases and amylases in their digestive tracts. A wide range of exoenzymes are secreted by *Bacillus* to complement the activity of the fish and to enhance its ability to digest nutrients enzymatically. Vadstein et al. (2018) have found that bacteria isolated from the digestive tracts of aquatic animals produce chitinases, proteases, cellulases, lipases, and trypsins.

### Improving the quality of the water

Probiotic bacteria directly absorb or decompose organic matter, improving water quality. Probiotics maintain a good water quality environment by decomposing fish or prawn excreta, food materials, plankton remains, and other organic matter by releasing CO<sub>2</sub>, nitrates, and phosphates. In culture water, Nimrat et al. (2012) found that mixed *Bacillus* probiotics significantly improved white shrimp's nitrite, pH, and ammonia levels. Kewcharoen and Srisapoom (2019) found that total ammonia was reduced when the probiotic *Bacillus subtilis* was added to shrimp culture water at 103-105 CFU/ml. At 106 CFU/ml, *Bacillus cereus* and *Pediococcus acidilactici* probiotics significantly decreased nitrate, ammonia, and biological oxygen demand in pond water (Khademzade et al., 2020). Several studies have shown that probiotic strains improve water quality, especially those of the gram-positive genus *Bacillus*. Converting organic matter into CO<sub>2</sub> makes this bacterial group more efficient than gram-negative bacteria. A high level of probiotics in production ponds can minimize the accumulation of dissolved and particulate organic carbon. Therefore, phytoplankton production can be balanced. El-Saadony et al., (2021). This hypothesis cannot be confirmed by cultivating shrimp or channel catfish with *Pseudomonas*, *Bacillus*, *Nitrobacter*, *Cellulomonas*, *Enterobacter*, or *Rhodopseudomonas*. Nitrifi-

cation is the only aspect of improving water quality that has received substantial attention (Pereira et al., 2022).

### Stress tolerance

Probiotics significantly improve stress tolerance in aquaculture species. Fish-fed *Lactobacillus plantarum* had a lower cortisol concentration than those fed control diets, which could be because cortisol is a stress hormone. Probiotics generally help fish to tolerate ammonia stress (Razafindralambo et al., 2023). *Nile tilapia* defense against hypoxia stress is significantly improved by *Aspergillus oryzae* probiotics (Dawood et al., 2019). Chowdhury et al. (2020) found that Pabda catfish supplemented with dietary commercial probiotics at 0.2 % were significantly more tolerant to saline water stress than those supplemented with higher and lower doses. As a result, probiotics-supplemented groups had lower plasma glucose levels, a stress indicator.

### Reproduction effects on aquatic species

The reproduction capacity of breeding aquaculture species depends on adequate lipids, proteins, fatty acids, vitamins C and E, and carotenoids. In addition, the interaction between these components affects reproduction in several ways, including fertility, fertilization, and larval development. Cultured fish species can be fed larger diets known as broodstock diets. Supplementing fresh fish with products from commercial diets improves the nutrition of fish hatcheries' broodstock. Fresh organisms used to feed broodstock fish include squid, cuttlefish, mussels, krill, and small crustaceans. Using unprocessed fish products often results in poor nutrition for broodstock fish, increasing the risk of parasites, bacteria, and viruses infecting the offspring and parents. Probiotics were added to food or water to prevent infections and study their effects on reproduction (Vidhya et al., 2019; Leong et al., 2023). An important study on the effects of probiotic supplements on fish reproductive performance was conducted by (Hasan et al., 2020), which included a strain of *B. subtilis* isolated from *Cirrhinus mrigala*'s intestine, incorporated at different concentrations into four ornamental fish species: *Poecilia reticulata*, *P. sphenops*, *Xiphophorus helleri*, and *X. maculatus*. Mixing *B. subtilis* at 106–108 cells per gram of food increased the gonadosomatic index, fecundity, viability, and larval production in all four species. Probiotics synthesize complex B vitamins, particularly thiamine (vitamin B1) and vitamin B12, which reduce the number of dead or deformed fry. Abasali et al. (2010) conducted a similar study using a commercial probiotic containing *Lactobacillus acidophilus*, *L. casei*, *Enterococcus faecium*, and *Bifidobacterium thermophilum*. Each female was evaluated for fry production and relative fecundity. Comparing the probiotic-treated group with the control group, 105 and 150 fry, on average, were found in the probiotic-treated group. In the

second study, 28 females were fecundated under control conditions, and 41 were fecundated under probiotic conditions.

### Enhancement of the immune response

In aquaculture species, probiotics can enhance a variety of immunological parameters. Through stimulation of the body's non-specific and cellular immunity, probiotics can inhibit pathogen infection (Hamka et al., 2020). By increasing the proportion of phagocytic cells in the head kidney and activating complement receptor expression, the administration of viable lactic acid bacteria, *Lactococcus lactis*, *Leuconostoc mesenteroides*, and *Lactobacillus sakei* improved both cellular and hormonal immune functions in rainbow trout. According to Doan et al., (2018), host-associated probiotics *Lactobacillus plantarum* and *Bacillus velezensis* improve skin mucus lysozyme and peroxidase activity, serum lysozyme and peroxidase activity, alternative complement, phagocytosis, and respiratory burst activity in *Nile tilapia*. The administration of *Bacillus pumilus*, a host gut-derived probiotic, to juvenile golden pompano, *Trachinotus ovatus* increased their lysozyme activity and total protein levels (Liu et al., 2019).

### Improve feed utilization

The use of probiotics alters enzymes and improves feed utilization, according to much research. *Lactobacillus pentosus* supplementation improved feed utilization in white shrimp, *Litopenaeus vannamei* (Zheng and Wang, 2016). *Nile tilapia* were shown to exhibit superior amylase, lipase, and protease activity after consuming heat-killed *L. plantarum* for 12 weeks (Dawood et al., 2019). *Astacus leptodactylus* fed with *L. plantarum* at concentrations of 107, 108, and 109 CFU/gm had elevated amylase, alkaline phosphatase, protease, and lipase activities (Valipour et al., 2019).

### Conclusion

Probiotics have been seen to positively impact the host's gut defense system, an important factor for preventing disease and treating conditions linked to inflammation within the digestive system. Probiotics have significantly improved aquatic animals' development performance and feed consumption by activating their digestive enzymes. Aquaculture species can benefit from probiotics in terms of their immunological characteristics. Boosting the host's immune system and promoting both cellular and non-specific immunity in the body, probiotics help prevent the spread of disease. As a result of their ability to absorb or break down harmful substances or organic matter, probiotic bacteria help clean water. As a result of their ability to break down organic debris like leftover food items, plankton remnants, and fish or prawn excreta, probiotics contribute

to maintaining a healthy water quality environment for cultured animals by assisting the nutrition cycle. As a substitute for pathogen inhibition and disease management in aquaculture species, probiotics are an effective alternative. Probiotic microorganisms can act as a barrier against opportunistic infections by releasing chemicals that kill or inhibit pathogenic bacteria in the host's intestine. There is a lot of pressure on crop species regarding aquaculture, which requires a high output level within a short period. The purpose of administering probiotics to food and water is to prevent illness and investigate their effects on reproduction. The probiotic supplements have a profound impact on fish reproduction.

### ACKNOWLEDGEMENTS

The authors thank the faculty members and researchers of Darrang College, Tezpur, Assam, India, and the College of Veterinary Science, Assam Agricultural University, Khanapara, Assam, India, for their encouragement and support. The authors are grateful to the DBT-ALSBT Hub (Assam) for providing micro-grants to Young Scientists of NER for the project on metagenomic analysis of the skin and gut microbiome of healthy and diseased fish.

### Conflict of Interest

The authors declare that they have no conflict of interest.

### REFERENCES

1. Abasali, H. & Mohamad, S. (2010). Effect of dietary supplementation with probiotic on reproductive performance of female livebearing ornamental fish. *Research Journal of Animal Sciences*, 4(4), 103-107.
2. Afrilasari, W. & Meryandini, A. (2016). Effect of probiotic *Bacillus megaterium* PTB 1.4 on the population of intestinal microflora, digestive enzyme activity and the growth of catfish (*Clarias* sp.). *HAYATI Journal of Biosciences*, 23 (4), 168-172.
3. Allameh, S.K., Noaman, V. & Nahavandi, R. (2017). Effects of probiotic bacteria on fish performance. *Adv Tech in Clin Microbiol* 1(2):11: <http://www.imedpub.com/advanced-techniques-in-clinicalmicrobiology>
4. Aly, S. M. (2010). Probiotics and aquaculture. *CABI Reviews*, (2009), 1-16.
5. Anee, I. J., Alam, S., Begum, R. A., Shahjahan, R. M. & Khandaker, A. M. (2021). The role of probiotics on animal health and nutrition. *The Journal of Basic and Applied Zoology*, 82, 1-16.
6. Armandeh, M., Mahmoudi, N. & Fallah Nosratabad, A. R. (2022). Screening and evaluation of phosphate-solubilizing bacteria isolated from aquaculture ponds in a step-by-step strategy as potential biofertilizer. *Journal of Applied Microbiology*, 133(3), 1581-1596.
7. Askarian F, Zhou Z, Olsen RE, Sperstad S. & Ringø E (2012) Culturable autochthonous gut bacteria in Atlantic

- salmon (*Salmo salar* L.) fed diets with or without chitin. Characterization by 16S rRNA gene sequencing, ability to produce enzymes and *in vitro* growth inhibition of four pathogens. *Aquaculture*, 1-8: 326-329.
8. Bahi, A., Guardiola, F. A., Messina, C. M., Mahdhi, A., Cerezuela, R., Santulli, A., ... & Esteban, M. A. (2017). Effects of dietary administration of fenugreek seeds, alone or in combination with probiotics, on growth performance parameters, humoral immune response and gene expression of gilthead seabream (*Sparus aurata* L.). *Fish & Shellfish Immunology*, 60, 50-58.
  9. Batista, S., Ramos, M. A., Cunha, S., Barros, R., Cristóvão, B., Rema, P., ... & Ozório, R. O. A. (2015). Immune responses and gut morphology of *Senegalese sole* (*Solea senegalensis*, K aup 1858) fed monospecies and multispecies probiotics. *Aquaculture Nutrition*, 21(5), 625-634.
  10. Bhatnagar, Anita & Saluja, Sonal. (2019). Synergistic effects of autochthonous probiotic bacterium and Mentha piperita diets in Catla catla (Hamilton, 1822) for enhanced growth and immune response. *Fisheries and Aquatic Sciences*. 22. 10.1186/s41240-019-0130-7.
  11. Butt, U. D., Lin, N., Akhter, N., Siddiqui, T., Li, S. & Wu, B. (2021). Overview of the latest developments in the role of probiotics, prebiotics, and synbiotics in shrimp aquaculture. *Fish & Shellfish Immunology*, 114, 263– 281.
  12. Cano-Lozano, J. A., Villamil Diaz, L. M., Melo Bolivar, J. F., Hume, M. E. & Ruiz Pardo, R. Y. (2022). Probiotics in tilapia (*Oreochromis niloticus*) culture: Potential probiotic Lactococcus lactis culture conditions. *Journal of Bioscience and Bioengineering*, 133(3), 187–194.
  13. Chowdhury, Gourab & Jinia, Masuma & Hossain, Md., Dey, Tanwi Islam, Md Jakiul Iqbal, Mohammed Akhtar, Shakil & Das, Bipresh. (2020). Effects of dietary probiotics on the growth, blood chemistry and stress response of Pabda catfish (*Ompok pabda*) juveniles. *AACL Bioflux*. 13. 1595-1605.
  14. Contreras□Tapia, R. A., Castellanos□Páez, M. E., Nandini, S., Benitez□Díaz□Mirón, M. I. & Garza□Mouriño, G. (2020). Enhancing rotifer *Brachionus calyciflorus* population growth with commercial probiotics. *Aquaculture research*, 51(5), 2102-2111.
  15. Coulibaly, W. H., Kouadio, N. G. R., Camara, F., Diguță, C. & Matei, F. (2023). Functional properties of lactic acid bacteria isolated from Tilapia (*Oreochromis niloticus*) in Ivory Coast. *BMC Microbiology*, 23(1), 152.
  16. Das, Susmita & Mondal, Kausik & Haque, Salma. (2017). A review on application of probiotic, prebiotic and synbiotic for sustainable development of aquaculture. 422-429
  17. Dawood, M. A., Koshio, S., Abdel□Daim, M. M., & Van Doan, H. (2019). Probiotic application for sustainable aquaculture. *Reviews in Aquaculture*, 11(3), 907-924.
  18. Dawood, M. A., Koshio, S., Ishikawa, M., & Yokoyama, S. (2015). Effects of partial substitution of fish meal by soybean meal with or without heat-killed Lactobacillus plantarum (LP20) on growth performance, digestibility, and immune response of amberjack, *Seriola dumerili* juveniles. *BioMed research international*, 2015.
  19. Dawood, M. A., Koshio, S., Ishikawa, M., El-Sabagh, M., Esteban, M. A., & Zaineldin, A. I. (2016). Probiotics as an environment-friendly approach to enhance red sea bream, *Pagrus major* growth, immune response and oxidative status. *Fish & Shellfish Immunology*, 57, 170-178.
  20. Dawood, Mahmoud, Magouz, Fawzy, Salem, Mahmoud & Abdel-Daim, Hanaa (2019). Modulation of digestive enzyme activity, blood health, oxidative responses and growth-related gene expression in GIFT byheat-killed *Lactobacillus plantarum* (L-137). *Aquaculture*. 505. 10.1016/j.aquaculture.2019.02.053.
  21. Dennis, E., & Uchenna, O. (2016). Use of probiotics as first feed of larval African catfish *Clarias gariepinus* (Burchell 1822). *Annual Research & Review in Biology*, 9 (2), 1-9.
  22. Doan, Hien & Hoseinifar, Seyed Hossein & Khanongnuch, Chartchai & Kanpiengjai, Apinun & Unban, Kridsada & Kim, Van & Srichaiyo, Satawat. (2018). Host-associated probiotics boosted mucosal and serum immunity, disease resistance and growth performance of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*. 491. 10.1016/j.aquaculture.2018.03.019.
  23. Dohail, A., Abdullah, M., Roshada, H. & Aliyu, M. (2009). Effects of the probiotic, Lactobacillus acidophilus, on the growth performance, hematology parameters and immunoglobulin concentration in African Catfish (*Clarias gariepinus*, Burchell 1822) fingerling. *Aquac Research*, 40 (14):1642-1652
  24. Duan, Y., Dong, H., Wang, Y., Zhang, Y. & Zhang, J. (2018). Effects of the dietary probiotic Clostridium butyricum on intestine digestive and metabolic capacities, SCFA content and body composition in *Marsupenaeus japonicus*. *Journal Ocean University China*, 17(3), 690-696
  25. Dysin, A. P., Egorov, A. R., Godzishhevskaya, A. A., Kirichuk, A. A., Tskhovrebov, A. G., & Kritchenkov, A. S. (2023). Biologically active supplements affecting producer microorganisms in food biotechnology: A review. *Molecules*, 28(3), 1413.
  26. El-Kady, A. A., Magouz, F. I., Mahmoud, S. A., & Abdel-Rahim, M. M. (2022). The effects of some commercial probiotics as water additive on water quality, fish performance, blood biochemical parameters, expression of growth and immune-related genes, and histology of Nile tilapia (*Oreochromis niloticus*). *Aquaculture*, 546, 737249.
  27. El-Saadony, M. T., Alagawany, M., Patra, A. K., Kar, I., Tiwari, R., Dawood, M. A., ... & Abdel-Latif, H. M. (2021). The functionality of probiotics in aquaculture: An overview. *Fish & shellfish immunology*, 117, 36-52.
  28. Gao, X. Y., Liu, Y., Miao, L. L., Li, E. W., Hou, T. T., & Liu, Z. P. (2017). Mechanism of anti-Vibrio activity of marine probiotic strain *Bacillus pumilus* H2, and characterization of the active substance. *AMB express*, 7, 1-10.
  29. Gobi, N., Malaikozhundan, B., Sekar, V., Shanthi, S., Vaseeharan, B., Jayakumar, R., & Nazar, A. K. (2016). GFP tagged *Vibrio parahaemolyticus* Dahv2 infection and the protective effects of the probiotic *Bacillus licheniformis* Dahb1 on the growth, immune and antioxidant responses in *Pangasius hypophthalmus*. *Fish & Shellfish Immunology*, 52, 230-238.
  30. Gupta, A., Gupta, P., & Dhawan, A. (2016). *Paenibacillus polymyxa* as a water additive improved immune response of *Cyprinus carpio* and disease resistance against *Aeromonas hydrophila*. *Aquaculture Reports*, 4, 86-92.
  31. Gutiérrez Ramirez, L. A., David Ruales, C. A., Montoya Campuzano, O. I., & Betancur Gonzalez, E. (2016). Efecto de la inclusión en la dieta de probióticos microencapsulados sobre algunos parámetros zootécnicos en ale-

- vinos de tilapia roja (*Oreochromis* sp.). *Revista de Salud Animal*, 38(2), 112-119.
32. Hai, N. V. (2015). The use of probiotics in aquaculture. *Journal of Applied Microbiology*, 119(4), 917-935.
  33. Hamka, Muhammad Subhan & Meryandini, Anja & Widanarni, Widanarni. (2020). Growth performance and immune response of catfish *Clarias* sp. given probiotics *Bacillus megaterium* PTB 1.4 and *Pediococcus pentosaceus* E2211. *Jurnal Akuakultur Indonesia*, 19. 50-60. 10.19027/jai.19.1.50-60.
  34. Hari Krishnan, R., Balasundaram, C. & Heo, M.S. (2010). *Lactobacillus sakei* BK19 enriched diet enhances the immunity status and disease resistance to streptococcosis infection in kelp grouper, *Epinephelus bruneus*. *Fish Shellfish Immunology*, 29(6):1037-1043
  35. Hasan, K. N. & Banerjee, G. (2020). Recent studies on probiotics as beneficial mediator in aquaculture: a review. *The Journal of Basic and Applied Zoology*, 81(1), 1-16.
  36. Hasan, K. N. & Banerjee, G. (2020). Recent studies on probiotics as beneficial mediator in aquaculture: a review. *The Journal of Basic and Applied Zoology*, 81, 1-16.
  37. Hoseinifar, S. H., Maradonna, F., Faheem, M., Hari Krishnan, R., Devi, G., Ringø, E., ... & Carnevali, O. (2023). Sustainable ornamental fish aquaculture: the implication of microbial feed additives. *Animals*, 13(10), 1583.
  38. Hoseinifar, S. H., Sun, Y. Z., Wang, A., & Zhou, Z. (2018). Probiotics as means of diseases control in aquaculture, a review of current knowledge and future perspectives. *Frontiers in microbiology*, 9, 2429.
  39. Ibrahim, M. D. (2015). Evolution of probiotics in aquatic world: Potential effects, the current status in Egypt and recent perspectives. *Journal of advanced research*, 6(6), 765-791.
  40. Iorizzo, M., Albanese, G., Letizia, F., Testa, B., Tremonte, P., Vergalito, F., Lombardi, S. J., Succi, M., Coppola, R., & Sorrentino, E. (2022). Probiotic Potentiality from Versatile *Lactiplantibacillus plantarum* Strains as Resource to Enhance Freshwater Fish Health. *Microorganisms*, 10(2), 463. <https://doi.org/10.3390/microorganisms10020463>
  41. Jha, Dilip & Bhujel, Ram & Anal, Anil. (2014). Dietary supplementation of probiotics improves survival and growth of Rohu (*Labeo rohita* Ham.) hatchlings and fry in outdoor tanks. *Aquaculture*. 10.1016/j.aquaculture.2014.10.026.
  42. Kaiser, D., Unger, D., Qiu, G., Zhou, H. & Gan, H. (2013). Natural and human influences on nutrient transport through a small subtropical Chinese estuary. *The Science of the Total Environment*, 450-451, 92-107.
  43. Karthiga Rani, M., Chelladurai, G. & Jayanthi, G. (2016). Isolation and identification of bacteria from marine market fish *Scomberomorus guttatus* (Bloch and Schneider, 1801) from Madurai district, Tamil Nadu, India. *Journal Parasit Disease*, 40(3), 1062-1065.
  44. Kewcharoen, W., & Srisapome, P. (2019). Probiotic effects of *Bacillus* spp. from Pacific white shrimp (*Litopenaeus vannamei*) on water quality and shrimp growth, immune responses, and resistance to *Vibrio parahaemolyticus* (AHPND strains). *Fish & Shellfish Immunology*, 94, 175-189.
  45. Khademzade, O., Zakeri, M., Haghi, M., & Mousavi, S. M. (2020). The effects of water additive *Bacillus cereus* and *Pediococcus acidilactici* on water quality, growth performances, economic benefits, immunohematology and bacterial flora of whiteleg shrimp (*Penaeus vannamei* Boone, 1931) reared in earthen ponds. *Aquaculture research*, 51(5), 1759-1770.
  46. Kiron V (2012) Fish immune system and its nutritional modulation for preventive health care. *Anim Feed Sci Technol.*, 173: 111-133.
  47. Knobloch, S., Skírnisdóttir, S., Dubois, M., Kolypczuk, L., Leroi, F., Leeper, A., ... & Marteinson, V. P. (2022). Impact of putative probiotics on growth, behavior, and the gut microbiome of farmed Arctic char (*Salvelinus alpinus*). *Frontiers in Microbiology*, 13, 912473.
  48. Lapatra, S. E., Fehringer, T. R., & Cain, K. D. (2014). A probiotic *Enterobacter* sp. provides significant protection against *Flavobacterium psychrophilum* in rainbow trout (*Oncorhynchus mykiss*) after injection by two different routes. *Aquaculture*, 433, 361-366.
  49. LaPatra, Scott, Fehringer, Tyson & Cain, Kenneth (2014). A probiotic *Enterobacter* sp. provides significant protection against *Flavobacterium psychrophilum* in rainbow trout (*Oncorhynchus mykiss*) after injection by two different routes. *Aquaculture*, 433. 361-366. 10.1016/j.aquaculture.2014.06.022.
  50. Lee, S., Katya, K., Park, Y., Won, S., Seong, M. & Bai, S. C. (2017). Comparative evaluation of dietary probiotics *Bacillus subtilis* WB60 and *Lactobacillus plantarum* KCTC3928 on the growth performance, immunological parameters, gut morphology and disease resistance in Japanese eel, *Anguilla japonica*. *Fish & Shellfish Immunology*, 61, 201-210.
  51. Leong, S. S., Korel, F., Lingoh, A. D., Sarbini, S. R., Toh, S. C., Abit, L. Y., & Wong, S. C. (2023). Current probiotics application for aquaculture feed: a review. *Borneo Science| The Journal of Science and Technology*, 44(2).
  52. Leyton, Y., Sayes, C., Mejias, C., Abarca, M., Wilson, R., & Senra, C. R. (2017). Increased larval survival of *Seriola lalandi* using *Pseudoalteromonas* sp. as probiotics. *Revista de Biología Marina y oceanografía*, 52(1), 95-101.
  53. Lin, H. L., Shiu, Y. L., Chiu, C. S., Huang, S. L., & Liu, C. H. (2017). Screening probiotic candidates for a mixture of probiotics to enhance the growth performance, immunity, and disease resistance of Asian seabass, *Lateolabrax japonicus* (Bloch), against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 60, 474-482.
  54. Lin, L. Z., Zheng, Q. W., Wei, T., Zhang, Z. Q., Zhao, C. F., Zhong, H., ... & Guo, L. Q. (2020). Isolation and characterization of fengycins produced by *Bacillus amyloliquefaciens* JFL21 and its broad-spectrum antimicrobial potential against multidrug-resistant foodborne pathogens. *Frontiers in Microbiology*, 11, 579621.
  55. Liu, Shubin, Shifeng, Wang Cai, Yan & Li, Er-Chao, Ren, Zhuling, Wu, Yue, Guo, Weiliang Sun, Yun & Zhou, Yongcan. (2019). Beneficial effects of a host gut-derived probiotic, *Bacillus pumilus*, on the growth, non-specific immune response and disease resistance of juvenile golden pompano, *Trachinotus ovatus*. *Aquaculture*. 514. 734446. 10.1016/j.aquaculture.2019.734446.
  56. Loh, J. Y. (2017). The role of probiotics and their mechanisms of action: an aquaculture perspective. *World Aquac*, 48, 19-23. Banerjee, G., Ray, A.K. & Askarian, F. (2013). Characterization and identification of enzyme

- producing autochthonous bacteria from the gastrointestinal tract of two Indian air-breathing fish. *Benef Microbes*, 4(3):277-284
57. Luo, L., Xu, Q., Xu, W., Li, J., Wang, C. A., Wang, L. & Zhao, Z. (2020). Effect of *Bacillus megaterium*-coated diets on the growth, digestive enzyme activity, and intestinal microbial diversity of Songpu mirror carp *Cyprinus specularis* Songpu. *BioMed Research International*, 2020.
  58. Maniat, M., Salati, A. P., Zanguee, N., Mousavi, S. M., & Hoseinifar, S. H. (2023). Effects of dietary *Pediococcus acidilactici* and Isomaltooligosaccharide on growth performance, immunity, and antioxidant defense in juvenile common carp. *Aquaculture Nutrition*, 2023.
  59. Martinez C.P., Ibáñez, A.L., Monroy Hermosillo, O.A. & Ramirez Saad, H.C. (2012). Use of probiotics in aquaculture. *Internatonal Sch Research Notices*, 5(1):55-59
  60. Mata, M. T., Luza, M. F., & Riquelme, C. E. (2017). Production of diatom–bacteria biofilm isolated from *Seriola lalandi* cultures for aquaculture application. *Aquaculture Research*, 48(8), 4308-4320.
  61. Meidong, R., Doolgindachbaporn, S., Sakai, K., & Tongpim, S. (2017). Isolation and selection of lactic acid bacteria from Thai indigenous fermented foods for use as probiotics in tilapia fish *Oreochromis niloticus*. *Aquaculture, Aquarium, Conservation & Legislation*, 10(2), 455-463.
  62. Melo, Javier, Ruiz-Pardo, Ruth, Hume, Michael, Sidjabat, Hanna & Villamil-Diaz, Luisa. (2020). Probiotics for cultured freshwater fish. *Microbiology Australia*. 10.1071/MA20026.
  63. Meng, Y., Zhang, X., Zhang, Z., Li, J., Zheng, P., Li, J., ... & Lu, Y. (2023). Effects of Microorganisms on Growth Performance, Body Composition, Digestive Enzyme Activity, Intestinal Bacteria Flora and Antimicrobial Peptide (AMP) Content of Black Soldier Fly Larvae (*Hermetia illucens*). *Animals*, 13(17), 2722.
  64. Merrifield, D. L., Bradley, G., Baker, R. T. M., & Davies, S. J. (2010). Probiotic applications for rainbow trout (*Oncorhynchus mykiss* Walbaum) II. Effects on growth performance, feed utilization, intestinal microbiota and related health criteria postantibiotic treatment. *Aquaculture Nutrition*, 16(5), 496-503.
  65. Merrifield, D.L., Dimitroglou, A., Bradley, G., Baker, R.T.M. & Davies, S.J. (2010) Probiotic applications for rainbow trout (*Oncorhynchus mykiss*, Walbaum) I. Effects on growth performance, feed utilization, intestinal microbiota and related health criteria. *Aquac Nutr*. 16(5), 504-510
  66. Michael ET, Amos SO, Hussaini LT (2014) A Review on Probiotics Application in Aquaculture. *Fish Aquac*, J 5: 111.
  67. Mocanu, E. E., Savin, V., Popa, M. D. & Dima, F. M. (2022). The effect of probiotics on growth performance, haematological and biochemical profiles in *Siberian sturgeon* (*Acipenser baerii* Brandt, 1869). *Fishes*, 7(5), 239.
  68. Mohapatra S, Chakraborty T, Prusty AK, Das P, Paniprasad K, et al. (2012) Use of different microbial probiotics in the diet of Rohu, *Labeo rohita* fingerling: Effects on growth, nutrient digestibility and retention, digestive enzyme activities and intestinal microflora. *Aquaculture Nutrition* 18: 1-11.
  69. Monier, M. N., Kabary, H., Elfeky, A., Saadony, S., El-Hamed, N. N. A., Eissa, M. E., & Eissa, E. S. H. (2023). The effects of *Bacillus* species probiotics (*Bacillus subtilis* and *B. licheniformis*) on the water quality, immune responses, and resistance of whiteleg shrimp (*Litopenaeus vannamei*) against *Fusarium solani* infection. *Aquaculture International*, 31(6), 3437-3455.
  70. Muchlisin, Z. A., Nazir, M., Fadli, N., Adlim, M., Hendri, A., Khalil, M., & Siti-Azizah, M. N. (2017). Efficacy of commercial diets with varying levels of protein on growth performance, protein and lipid contents in carcass of *Acehnese mahseer*, *Tor tambra*. <http://hdl.handle.net/1834/12173>
  71. Mujeeb R.K.M., Jesmi, Y., Thomas, A.P. & Mohamed Hatha, A.A. (2010). Probiotic effect of *Bacillus* NL110 and *Vibrio* NE17 on the survival, growth performance and immune response of *Macrobrachium rosenbergii* (de Man). *Aquac Research*, 41(9), 120-134.
  72. Munirasu, S., Ramasubramanian, V., & Arunkumar, P. (2017). Effect of Probiotics diet on growth and biochemical performance of freshwater fish *Labeo rohita* fingerlings. *Journal of Entomology and Zoology Studies*, 5(3), 1374-1379.
  73. Muñoz-Atienza, E., Araújo, C., Magadán, S., Hernández, P. E., Herranz, C., Santos, Y., & Cintas, L. M. (2014). In vitro and in vivo evaluation of lactic acid bacteria of aquatic origin as probiotics for turbot (*Scophthalmus maximus* L.) farming. *Fish & Shellfish Immunology*, 41(2), 570-580.
  74. Nakano, T. (2007). Microorganism. En dietary supplements for the health and quality of cultured fish. *CA International, Lon-don, UK*.
  75. Nandi, A., Banerjee, G., Dan, S. K., Ghosh, K., & Ray, A. K. (2017). Probiotic efficiency of *Bacillus* sp. in *Labeo rohita* challenged by *Aeromonas hydrophila*: assessment of stress profile, haemato-biochemical parameters and immune responses. *Aquaculture research*, 48(8), 4334-4345.
  76. Nayak SK (2010) Probiotics and immunity: A fish perspective. *Fish Shellfish Immunol.*, 29: 2-14.
  77. Naylor, R.L., Hardy, R.W., Buschmann, A.H. et al. A 20-year retrospective review of global aquaculture. *Nature* 591, 551–563 (2021). <https://doi.org/10.1038/s41586-021-03308-6>
  78. Ng, W. K., Kim, Y. C., Romano, N., Koh, C. B., & Yang, S. Y. (2014). Effects of dietary probiotics on the growth and feeding efficiency of red hybrid tilapia, *Oreochromis* sp., and subsequent resistance to *Streptococcus agalactiae*. *Journal of Applied Aquaculture*, 26(1), 22-31.
  79. Nimrat, S., Boonthai, T. & Vuthiphandchai, V. (2011). Effects of probiotic forms, compositions of and mode of probiotic administration on rearing of Pacific white shrimp (*Litopenaeus vannamei*) larvae and postlarvae. *Animal Feed Science Technology*, 169(3-4):244-258
  80. Nimrat, Subuntith & Suksawat, Sunisa & Boonthai, Traimat & Vuthiphandchai, Verapong. (2012). Potential *Bacillus* probiotics enhance bacterial numbers, water quality and growth during early development of white shrimp (*Litopenaeus vannamei*). *Veterinary Microbiology*, 159. 443-50. 10.1016/j.vetmic.2012.04.029.
  81. Noshair, I., Kanwal, Z., Jabeen, G., Arshad, M., Yunus, F. U. N., Hafeez, R., ... & Alomar, S. Y. (2023). Assessment of dietary supplementation of *Lactobacillus rhamnosus* probiotic on growth performance and disease resistance in *Oreochromis niloticus*. *Microorganisms*, 11 (6), 1423.

82. Okeke, E.S., Chukwudozie, K.I., Nyaruaba, R. et al. Antibiotic resistance in aquaculture and aquatic organisms: a review of current nanotechnology applications for sustainable management. *Environ Sci Pollut Res* 29, 69241–69274 (2022). <https://doi.org/10.1007/s11356-022-22319-y>
83. Omenwa, V.C., Mbakwem-Aniebo, C. & Ibiene, A.A. (2015). Effects of selected probiotics on the growth and survival of fry-fingerlings of *Clarias gariepinus*. *Pharm Biology Science*, 10:89-93.
84. Pannu, & Dahiya, & Sabhlok, & Kumar, Suresh & Sarsar, Vikas & Gahlawat, S.. (2014). Effect of probiotics, antibiotics and herbal extracts against fish bacterial pathogens. *Ecotoxicol. Environ. Contam.*, 9. 13-20.
85. Park, Y., Lee, S., Hong, J., Kim, D., Moniruzzaman, M., & Bai, S. C. (2017). Use of probiotics to enhance growth, stimulate immunity and confer disease resistance to *Aeromonas salmonicida* in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Research*, 48(6), 2672-2682.
86. Pegu, A., Kalita, R., Das, P. & Baruah, C. (2023). Nutritional composition of small indigenous species of fishes of Northeast India: A systematic review. *Journal of Applied and Natural Science*, 15(2), 649 -662. <https://doi.org/10.31018/jans.v15i2.4490>
87. Perdichizzi, A., Meola, M., Caccamo, L., Caruso, G., Gai, F., & Maricchiolo, G. (2023). Live Yeast (*Saccharomyces cerevisiae* var. *boulardii*) Supplementation in a European Sea Bass (*Dicentrarchus labrax*) Diet: Effects on the Growth and Immune Response Parameters. *Animals*, 13 (21), 3383.
88. Pereira, W. A., Mendonça, C. M. N., Urquiza, A. V., Marteinsson, V. P., LeBlanc, J. G., Cotter, P. D., & Oliveira, R. P. (2022). Use of probiotic bacteria and bacteriocins as an alternative to antibiotics in aquaculture. *Microorganisms*, 10(9), 1705.
89. Pereira, W. A., Mendonça, C. M. N., Urquiza, A. V., Marteinsson, V. P., LeBlanc, J. G., Cotter, P. D., & Oliveira, R. P. (2022). Use of probiotic bacteria and bacteriocins as an alternative to antibiotics in aquaculture. *Microorganisms*, 10(9), 1705.
90. Razafindralambo, H., Vijayaram, S., Rabetafika, H., Nguyen, H. T., & Sun, Y. Z. (2023). Probiotics in Aquaculture. *Encyclopedia*.
91. Reque, P. M., & Brandelli, A. (2022). An introduction to probiotics. In *Probiotics* (pp. 1-17). Academic Press.
92. Ringo E, Sperstad S, Myklebust R, Refstie S, Krogdahl Å (2006) Characterization of the microbiota associated with intestine of Atlantic cod (*Gadus morhua* L.) The effect of fish meal, standard soybean meal and abioprocessed soybean meal. *Aquaculture* 261: 829-841.
93. Rodriguez-Estrada, U., Satoh, S., Haga, Y., Fushimi, H., & Sweetman, J. (2013). Effects of inactivated *Enterococcus faecalis* and mannan oligosaccharide and their combination on growth, immunity, and disease protection in rainbow trout. *North American Journal of Aquaculture*, 75(3), 416-428.
94. Romarheim, O.H., Øverland, M., Mydland, L.T., Skrede, A. & Landsverk, T. (2011). Bacteria grown on natural gas prevent soybean meal-induced enteritis in Atlantic salmon. *Journal Nutrition*, 141(1):124-130.
95. Safari, R., Adel, M., Lazado, C. C., Caipang, C. M. A., & Dadar, M. (2016). Host-derived probiotics *Enterococcus casseliflavus* improves resistance against *Streptococcus iniae* infection in rainbow trout (*Oncorhynchus mykiss*) via immunomodulation. *Fish & Shellfish Immunology*, 52, 198 -205.
96. Sahandi, Javad & Jafaryan, Hojatollah & Soltani, Mehdi & Ebrahimi, Pouneh. (2018). The Use of Two Bifidobacterium Strains Enhanced Growth Performance and Nutrient Utilization of Rainbow Trout (*Oncorhynchus mykiss*) Fry. *Probiotics and Antimicrobial Proteins*, 11. 10.1007/s12602-018-9455-2.
97. Sanches-Fernandes, G. M., Sá-Correia, I., & Costa, R. (2022). Vibriosis outbreaks in aquaculture: addressing environmental and public health concerns and preventive therapies using gilthead seabream farming as a model system. *Frontiers in Microbiology*, 13, 904815.
98. Sarker, P. K. (2023). Microorganisms in fish feeds, technological innovations, and key strategies for sustainable aquaculture. *Microorganisms*, 11(2), 439.
99. Sayes, C., Leyton, Y., & Riquelme, C. E. (2016). Bacterium *Pseudoalteromonas* sp. potential probiotic for larval fish culture. *Latin American Journal of Aquatic Research*, 44(1), 76-84.
100. Schaeck, Marlien & Reyes-López, Felipe & Vallejos-Vidal, Eva & Van Cleemput, Jolien & Duchateau, Luc & Van den Broeck, Wim & Tort, Lluís & Decostere, Annemie. (2017). Cellular and transcriptomic response to treatment with the probiotic candidate *Vibrio lentus* in gnotobiotic sea bass (*Dicentrarchus labrax*) larvae. *Fish & Shellfish Immunology*, 63. 10.1016/j.fsi.2017.01.028
101. Sharifuzzaman, S. M., Rahman, H., Austin, D. A., & Austin, B. (2018). Properties of probiotics *Kocuria* SM1 and *Rhodococcus* SM2 isolated from fish guts. *Probiotics and Antimicrobial Proteins*, 10, 534-542.
102. Shewale, R.N. & Sawale, Pravin & Khedkar, Chandraprakash & Singh, Ajay. (2014). Selection criteria for probiotics: A review. *International Journal of Probiotics and Prebiotics*, 9. 17-22.
103. Silarudee, Somrudee & Tongpim, Saowanit & Charoensri, Nicha & Doolgindachbaporn, Sompong. (2019). Effect of a Probiotic *Lactobacillus plantarum* CR1T5 Dietary on Non-specific Immunity in Black Eared Catfish (*Pangasius larnaudii*). *Journal of Pure and Applied Microbiology*, 13. 289-296. 10.22207/JPAM.13.1.31.
104. Sîrbu, E., Dima, M. F., Tenciu, M., Cretu, M., Coadă, M. T., Țoțoiu, A., & Patriche, N. (2022). Effects of dietary supplementation with probiotics and prebiotics on growth, physiological condition, and resistance to pathogens challenge in Nile tilapia (*Oreochromis niloticus*). *Fishes*, 7(5), 273.
105. Sîrbu, E., Dima, M. F., Tenciu, M., Cretu, M., Coadă, M. T., Țoțoiu, A., & Patriche, N. (2022). Effects of dietary supplementation with probiotics and prebiotics on growth, physiological condition, and resistance to pathogens challenge in Nile tilapia (*Oreochromis niloticus*). *Fishes*, 7(5), 273.
106. Sumon, M. A. A., Sumon, T. A., Hussain, M. A., Lee, S. J., Jang, W. J., Sharifuzzaman, S. M., & Hasan, M. T. (2022). Single and multi-strain probiotics supplementation in commercially prominent finfish aquaculture: Review of the current knowledge. *Journal of Microbiology and Biotechnology*, 32(6), 681.

107. Sunitha, K. & Krishna, P.V. (2016). Efficacy of probiotics in water quality and bacterial biochemical characterization of fish ponds. *International Journal Curr Microbiology Appl Science*, 5(9), 30-37
108. Taoka, Y., Maeda, H., Jo, J. Y., Jeon, M. J., Bai, S. C., Lee, W. J., & Koshio, S. (2006). Growth, stress tolerance and non-specific immune response of Japanese flounder *Paralichthys olivaceus* to probiotics in a closed recirculating system. *Fisheries Science*, 72, 310-321.
109. Taoka, Y., Maeda, H., Jo, J. Y., Kim, S. M., Park, S. I., Yoshikawa, T., & Sakata, T. (2006). Use of live and dead probiotic cells in tilapia *Oreochromis niloticus*. *Fisheries Science*, 72, 755-766.
110. Torres-Maravilla, E., Parra, M., Maisey, K., Vargas, R. A., Cabezas-Cruz, A., Gonzalez, A., & Bermúdez-Humarán, L. G. (2024). Importance of Probiotics in Fish Aquaculture: Towards the Identification and Design of Novel Probiotics. *Microorganisms*, 12(3), 626.
111. Tovar-Ramírez, D., Mazurais, D., Gatesoupe, J. F., Quazuguel, P., Cahu, C. L., & Zambonino-Infante, J. L. (2010). Dietary probiotic live yeast modulates antioxidant enzyme activities and gene expression of sea bass (*Dicentrarchus labrax*) larvae. *Aquaculture*, 300(1-4), 142-147.
112. Vadstein, O., Attramadal, K. J., Bakke, I., Forberg, T., Olsen, Y., Verdegem, M., ... & Bossier, P. (2018). Managing the microbial community of marine fish larvae: a holistic perspective for larviculture. *Frontiers in Microbiology*, 9, 1820.
113. Valipour, Alireza & Nedaei, Shiva & Noori, Ahmad & Khanipour, Ali & Hoseinifar, Seyed Hossein. (2019). Dietary Lactobacillus plantarum affected on some immune parameters, air-exposure stress response, intestinal microbiota, digestive enzyme activity and performance of narrow clawed Eschscholtz crayfish *Astacus leptodactylus*. 10.1016/j.aquaculture.2019.01.064. *leptodactylus*, 504.
114. Vidhya, H. S., Chandrasekaran, N., Mukherjee, A. & Thomas, J. (2019). Use of Probiotics as a Feed in Aquaculture: A Review. *Chemical Technology and Informatics in Chemistry with Applications*, 3-17.
115. Wang, Y., Sun, J., Zhong, H., Li, N., Xu, H., Zhu, Q., & Liu, Y. (2017). Effect of probiotics on the meat flavour and gut microbiota of chicken. *Scientific Reports*, 7(1), 6400.
116. Wanka, K.M., Damerau, T., Costas, B., Krueger, A., Schulz, C. & Wuertz, S. (2018). Isolation and characterization of native probiotics for fish farming. *Bmc Microbiology* 18(1):1-13
117. Wesseling, W., Lohmeyer, M., Wittka, S., Bartels, J., Kroll, S., Soltmann, C., & Meinhardt, F. (2016). Adverse effects of immobilised *Pseudoalteromonas* on the fish pathogenic *Vibrio anguillarum*: an in vitro study. *Journal of Marine Sciences*, 2016.
118. Xu, H.M., Rong, Y.J., Zhao, M.X., Song, B. & Chi, Z.M. (2014). Antibacterial activity of the lipopeptides produced by *Bacillus amyloliquefaciens* M1 against multidrug-resistant *Vibrio* spp. isolated from diseased marine animals. *App Microbiology Biotechnol.*, 98(1):127–36. 10.1007/s00253-013-5291-1
119. Yamashita, M. M., Pereira, S. A., Cardoso, L., De Araujo, A. P., Oda, C. E., Schmidt, É. C., & Mourinho, J. L. P. (2017). Probiotic dietary supplementation in Nile tilapia as prophylaxis against streptococcosis. *Aquaculture Nutrition*, 23(6), 1235-1243.
120. Yu, Y. M., Poon, P. M. Y., Sharma, A. A., Chan, S. M. N., Lee, F. W. F., Mo, I. W. Y., & Sze, E. T. P. (2022). Colonization of *Lactobacillus rhamnosus* GG in *Cirrhinus molitor* (mud carp) fingerling: evidence for improving disease resistance and growth performance. *Applied Microbiology*, 2(1), 175-184.
121. Zhang, S., Lin, L., & Wang, X. (2021). Optimization of a marine fish release strategy: a case study of black sea bream *Acanthopagrus schlegelii* in the Zhanjiang estuary, Northern south China Sea. *Frontiers in Environmental Science*, 9, 779544.
122. Zhao, J., Pan, J., Zhang, Z., Chen, Z., Mai, K., & Zhang, Y. (2023). Fishmeal Protein Replacement by Defatted and Full-Fat Black Soldier Fly Larvae Meal in Juvenile Turbot Diet: Effects on the Growth Performance and Intestinal Microbiota. *Aquaculture Nutrition*, 2023.
123. Zheng, Chen & Wang, Wei. (2016). Effects of *Lactobacillus pentosus* on the growth performance, digestive enzyme and disease resistance of white shrimp, *Litopenaeus vannamei* (Boone, 1931). *Aquaculture Research*. 48. 10.1111/are.13110.
124. Ziaei-Nejad, S., Rezaei, M. H., Takami, G. A., Lovett, D. L., Mirvaghefi, A. R., & Shakouri, M. (2006). The effect of *Bacillus* spp. bacteria used as probiotics on digestive enzyme activity, survival and growth in the Indian white shrimp *Fenneropenaeus indicus*, *Aquaculture*, 252(2-4), 516-524.