



Mutagenic effectiveness and efficiency of gamma rays in snake gourd (*Trichosanthes anguina* L.)

P. Sidhya* and M.K. Pandit

Department of Vegetable crops, Faculty of Horticulture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur-741252 (W.B.), INDIA

*Corresponding author. E-mail: palash.sum@gmail.com

Received: February 14, 2015; Revised received: July 12, 2015; Accepted: August 14, 2015

Abstract: The present experiment was undertaken to determine the mutagenic effectiveness and efficiency of gamma rays on different biological parameters in snake gourd. The research was conducted in two generations namely M₁ and M₂ during spring-summer season (mid- February) of 2012 and 2013 at the Horticulture research station, Mondoury, Bidhan Chandra Krishi Viswavidyalaya. The parent material, selfed seeds of BCSG-28 variety were irradiated with 100 Gy, 150 Gy, 200 Gy, 250 Gy and 300 Gy doses of gamma rays. The seeds along with control were space planted for raising M₁ generation. Each M₁ plant was harvested separately and desirable M₁ individual plant progeny rows were laid in RBD for raising M₂ generation. The effectiveness and efficiency of the mutagen used was assessed from the data on biological damage in M₁ generation. In M₁ generation, results showed a dose dependent retardation in biological parameters like seed germination, plant survival and 200 Gy was depicted as LD₅₀ indicating less damaging effect at lower doses on genetic material. In M₂ generation, Lower doses 100 Gy (28.80 effectiveness, 21.58 efficiency) and 150 Gy (18.33 effectiveness, 8.68 efficiency) treatments were found as effective and efficient and a wide range of induced variability was observed in almost all traits. The mutants with short fruit, higher fruit diameter and reduced vine length were isolated in M₂ generation.

Keywords: Gamma rays, Grey (Gy), Mutagenic effectiveness, Mutagenic efficiency, Snake gourd

INTRODUCTION

Snake gourd (*Trichosanthes anguina* L.), a monoecious minor cucurbit having 2n=22, plays an important role in human diet. Snake gourd is a commonly grown vegetable in the country, originated in Indian Archipelago and widely distributed in humid tropical areas of many countries, including Southeast Asia, China, Japan, Australia, Fiji, Mauritius, Java, South America and some parts of America. In India it is cultivated in Kerala, Tamilnadu, Meghalaya, Tripura, Assam, West Bengal and Odissa. Flowers are white in colour and fringed. The fruits are long narrow, cylindrical, slender, long tapering and may vary from 1-6 feet in length. It is important as a good source of minerals, fibre and nutrients to make the food wholesome and healthy (Ahmed *et al.*, 2000).

Mutations occur in natural populations spontaneously at a low rate. Mutation offer a great scope in breeding any crop and can be incorporated into cropping programme as conventional allow obtaining genotypes with desirable traits (Ahloowalia *et al.*, 2004). So, induction of mutation through radiation or chemicals are now very common worldwide to boost up crop wealth and food security. The FAO estimates that about 800 million people do not have enough food to eat. Global food security continues to be the central issue to cope up

with the ever growing human population (Kharkwal and Shu 2009). Sheeba *et al.*, (2005) reported that in *Sesamum indicum* L. seed germination, seedling survival, plant height and pollen fertility were reduced significantly with the increase in dosage levels of both gamma rays and EMS. Adamu *et al.* (2002) also recorded decreased seedling emergence, seedling height and root length, and seedling survival in groundnut with the treatment of gamma rays. Kumar and Yadav (2010) had reported significant plant height reduction in sesame due to higher doses of mutagenic treatment. The efficacy of any mutagen in plant breeding depends on its effectiveness and efficiency (Konzak *et al.*, 1965). Induced genetic variability is essential for any crop improvement programme and its creation and management are central to plant breeding. The present investigation was carried out to determine LD₅₀ in M₁ generation and to assess the efficiency and effectiveness of gamma ray mutagen in M₂ generation in snake gourd.

MATERIALS AND METHODS

The parent materials used in the present experiment was selfed seeds of BCSG-28, a local variety of snake gourd with green rind colour with distinct white stripes and smooth surface, elongated, 50-55cm in length and 3.5-4.5 cm in girth. Uniform, healthy and dry seeds of

snake gourd were irradiated with different doses viz. 100, 150, 200, 250 and 300 Gy (Grey) of gamma rays (source: ^{60}Co) Jadavpur technical campus, Department of Atomic Energy, Kolkata. Next day morning, treated seeds of each dose and control were sown during spring-summer (mid-February) 2012 for raising M_1 generation.

Seed germination was calculated as proportion of seeds germinated in each treatment including control and it was recorded 25 days after sowing in the field. Plant survival was calculated as proportion of plants survived until maturity in each treatment including control. LD_{50} was depicted by drawing graph of plant survival % versus dosage of gamma rays. In M_2 generation, M_1 plant progeny rows were raised and mutants were observed.

Assessment of mutation frequency, mutagenic effectiveness and efficiency: Individual plants of the treatments including control were harvested separately in treatment wise and desirable M_1 mutants were raised in M_2 generation; then mutation frequency, Mutagenic effectiveness and efficiency were calculated (Konzak *et al.*, 1965) and expressed in percentage.

Mutation frequency: On M_2 plant basis (% of mutated M_2 plants) Mutation frequency was estimated as per cent of segregating M_1 Plant progenies (Gaul, 1964).

Mutagenic effectiveness: The ratio of "factor mutations to doses" means frequency of a given mutation at a given dose of mutagen employed. Mutagenic effectiveness was calculated by the formula :

$$\text{Mutagenic Effectiveness} = \frac{\text{Mutation frequency in } M_2}{\text{Dose of mutagen (kR)}}$$

Where, Gy=Grey, 10 Grey= 1 kR (kilo Roentgen of gamma rays)

Mutagenic efficiency: The ratio of "Factor mutations to biological damage" means desirable changes free from associated undesirable changes on mutagenesis. Mutagenic efficiency was calculated by the formula:

$$\text{Mutagenic Efficiency} = \frac{\text{Mutation frequency in } M_2}{\% \text{ lethality or sterility in } M_1}$$

Where, Lethality was calculated as percentage of plant survival reduction and sterility was calculated as per cent of seed fertility reduction.

RESULTS AND DISCUSSION

Data on germination and plant survival are presented in Table-1. In M_1 generation dose dependent retardation in seed germination was observed. As compared to control maximum seed germination was recorded 93.33 per cent, followed by 81.66 per cent in the 100 Gy and 150 Gy treatments respectively while minimum seed germination was recorded to be 36.66 per cent at highest dose 300 Gy (Table-1). Similar results have been observed by Kumar and Yadav (2010). It was observed that increase in radiation dose, reduced germination percentage, which was also recorded by Ciftci *et al.*, (2006) and Kumari and Singh (1996). Similarly seed germination percentage and plant

survival also decreased with increased dose of γ -ray in the present experiment. It indicates that significant increase in lethality takes place at higher doses of mutagen. The similar results have been revealed by Vinod and Sharma (1998). The dose 200 Gy was found out to be the LD_{50} as it recorded fifty per cent plant survival (Table-1). The same trend was also noted by Shevchenko *et al.*, (1990) in pea and Omar *et al.*, (2008) in chilli. Genetic variability has been reported to increase in doses of gamma rays (Gupta, 1970)

The mutagenic effectiveness is a measure of point mutations induced by a unit dose of mutagen while mutagenic efficiency represents the proportion of mutation in relation to the associated undesirable biological effects, such as chromosomal aberration, lethality and sterility induced by mutagen (Konzak *et al.*, 1965). In fact the ratio of chlorophyll mutations induced in M_2 generation to the various biological damages induced in M_1 generation such as lethality, and pollen sterility indicates efficiency of a mutagen. In M_2 generation, mutagenic effectiveness and efficiency of the mutagen exhibited inverse relationship with increase in dose (Table-2). Most effective (28.8 %) and efficient (21.58 %) gamma radiation dose was 100 Gy on the basis of mutant plants in M_2 generation. Lower doses of gamma rays recorded higher efficiency in the present study which is in tune with the findings of Paul and Singh (2005). Maximum number of mutated plants are observed in 100 Gy and 150 Gy (3) followed by 200 Gy (2). The highest mutation frequency was observed in 100 Gy (2.88) followed by 150 Gy (2.75). The low variability for efficiency can be interpreted by assuming that the extent of damage in M_1 generation determines the mutability of genes, irrespective of mutagen used. Similar results also recorded by Gautam *et al.*, (1992) and Sharma *et al.*, (2006) in Urdbean. The decrease in seed germination induced by mutagenic treatments may be the result of damage of cell constituents at molecular level or altered enzyme activity (Khan and Goyel, 2009; Chowdhury and Tah, 2011). Srivastava *et al.* (2011) suggested the reduction in seedling survival of wheat was due to the hindrance caused by the mutagen on different metabolic pathway of the cells.

Conclusion

The experimental results revealed maximum seed germination percentage (93.33) and plant survival percentage (86.66) of snake gourd, *Trichosanthes anguina* in 100 Gy after the control (100, 100 respectively). 200 Gy was noted LD_{50} and the lower doses (100 Gy and 150 Gy) are more effective and efficient. The mutants with short fruit, higher fruit diameter and reduced vine length were isolated in M_2 generation. Lower effective doses of gamma rays may contribute to considerable utility.

ACKNOWLEDGEMENTS

The authors are grateful to the Department of Vegetable Crops, Faculty of Horticulture, Bidhan Chandra

Table 1: Estimation of seed germination (%) and plant survival (%) in M₁ generation

S N.	Dose	Seed germination percentage (%)	Per cent reduction	Plant survival percentage (%)	Per cent reduction
1.	Control	100	00.00	100	00.00
2.	100 Gy	93.33	6.67	86.66	13.34
3.	150 Gy	81.66	18.34	68.33	31.67
4.	200 Gy	58.33	41.67	50.00	50.00
5.	250 Gy	43.33	56.67	38.33	61.67
6.	300 Gy	36.66	63.34	26.66	73.34

Table 2: Assessment of Mutagenic Effectiveness and Mutagenic Efficiency in M₂ generation.

S N.	Dose	M ₂ plant population	Number of M ₂ plants mutated	Mutation frequency (Mf)	Lethality or per cent plant survival reduction (L)	Mutagenic effectiveness Mf X 100/ Dose	Mutagenic efficiency Mf X 100/ L
1.	Control	-	-	-	-	-	-
2.	100 Gy	104	3	2.88	13.34	28.80	21.58
3.	150 Gy	109	3	2.75	31.67	18.33	8.68
4.	200 Gy	103	2	1.94	50.00	9.70	3.88
5.	250 Gy	102	1	0.98	61.67	3.92	1.58
6.	300 Gy	106	1	0.94	73.34	3.13	1.28

Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal, for providing the necessary research facilities.

REFERENCES

- Adamu, A.K. Oluranju, P.E. Bate, J.A. and Ogunlade, O.T. (2002). Radio sensitivity and effective dose determination in groundnut (*Arachis hypogaea* L.) irradiated with gamma-rays. *J. Agric. and Env*, 3(1): 17-84.
- Ahloowalia, B.S. Maluszynski, M. and Nichterlein, K. (2004). Global impact of mutation-derived varieties. *Euphytica*, 135: 187-204.
- Ahmed, M.S. Rasul, M.G. Bashir, M.K. and Mian, A.S.M. (2000). Variability and heterosis in snake gourd (*Trichosanthes anguina* L.). *Bangladesh J. Plant Breed. Genet*, 13: 27-32.
- Chowdhury, R. and Tah, J. (2011). Assessment of Chemical Mutagenic Effects in Mutation Breeding Programme for M₁ Generation of Carnation (*Dianthus caryophyllus*). *Research in Plant Biology*, 1(4): 23-32.
- Ciftci, C.Y. Turkan, A.D. Khawar, K.M. Atak, M. and Ozcan, S. (2006). Use of gamma rays to induce mutations in four pea cultivars. *Turkish Journal of Biology*, 30(1): 29-37.
- Gaul, H. (1964). Mutation in plant breeding. *Radiation Botany*, 2(1): 12-16.
- Gautam, A.S. Sood, K.C. and Richarria, A.K. (1992). Mutagenic Effectiveness and Efficiency of gamma rays, Ethyl methane sulphate and their synergistic effects in Black gram (*Vigna mungo* L.). *Cytologia*, 57: 85-89.
- Gupta, A. K. (1970). Induced polygenic variability in diploid and tetraploid barley. Radiation and radio-metric substances in mutation breeding (Proc. Symp. Bombay. 1969). Department of Atomic Energy, India, 324-327.
- Khan, S. and Goyal, S. (2009). Improvement of Mungbean Varieties through induced mutations. *Afr. J. Plant Sci*, 3: 174-180.
- Kharkwal, M.C. and Shu, Q.Y. (2009). The role of induced mutations in world food security. In: Induced plant mutations in the genomics era. Shu, Q.Y. (Ed.). *Food and Agriculture Organization of the United Nations, Rome, Italy*, pp 33-38.
- Konzak, C.F. Nilan, R.A. Wagner, J. and Foster, R.J. (1965). Efficient chemical mutagenesis. In: "The use of induced mutation in plant breeding", *Rad. Bot.* (Supl.), 5: 49-70.
- Kumar, G. and Yadav, R.S. (2010). Ems induced genomic disorders in sesame (*Sesamum indicum* L.). *Rom. J. Biol*, 55(2): 97-10.
- Kumari, R. and Singh, Y. (1996). Effect of gamma rays and EMS on seed germination and plant survival of *Pisum sativum* L. and *Lens Culinaris* Medic. *Neo Botanica*, 4 (1): 25-29.
- Omar, S.R. Ahmed, O. H. Saamin, S and Majid, N. M. (2008). Gamma Radio Sensitivity study on chili (*Capscium annum*), *American Journal of Applied Sciences*, 5(2) 67-70.
- Paul, A. and Singh, D.P. (2005). Frequency, spectrum and segregation pattern of chlorophyll and macro mutations in field pea (*pisum sativum*). *Journal of Legume Research*, 28(2) 94-98.
- Sharma, A.K. Singh, V.P. and Singh, R.M. (2006). Efficiency and Effectiveness of the gamma rays, EMS and their combination in Urd bean. *Indian Journal of Pulses Research*, 19(1): 111-112.
- Sheeba, A. Abumalarmalhi, J. Babu, S and Ibrahim, S.M. (2005). Mutagenic effects of gamma rays and EMS in M₁ generation in sesame. *Resource in crops*, 6(2): 300-306.
- Shevchenko, A.M. Denisenko, E.G. and Blyandur, O.V. (1990). Use of gamma rays in producing source material in breeding determinate pea varieties. *Sel. Skokhozyist Vennaya Radio Biologiya*, 4-12.
- Srivastava, P. Marker, S. Pandey, P. and Tiwari, D.K. (2011). Mutagenic Effects of Sodium Azide on the Growth and Yield Characteristics in Wheat (*Triticum aestivum* L. em. Thell.). *Asian J. Plant Sci*, 10: 190-201.
- Vinod and Sharma, B. (1998). Primary mutagenic damage and mutagenic effectiveness in dwarf peas. *Crop Research Hisar*, 16(3): 340-343.