



Response of different wheat varieties towards *Azospirillum* and phosphate solubilizing bacteria (PSB) seed inoculation

Animesh Pathak^{1*}, S. K. Chakrabarti², Rajib Das³ and M K. Mandal³

¹Department of Seed Science and Technology, M. S. Swaminathan School of Agriculture, Centurion University of Technology and Management, Paralakhemundi, Dist. Gajapati-761211 (Odisha), INDIA

²Department of Seed Science and Technology, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741252 (W.B.), INDIA

³Department of Agronomy, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia-741252 (W.B.), INDIA

*Corresponding author. E-mail: animeshpathak88@gmail.com

Received: May 29, 2015; Revised received: August 27, 2015; Accepted: February 15, 2016

Abstract: Present investigation was undertaken to utilize *Azospirillum* and Phosphate Solubilizing Bacteria (PSB) to avail atmospheric nitrogen and soluble phosphate in soil for the wheat crop. The sonalika variety (V_1) responded best towards seed inoculation by *Azospirillum* and Phosphate Solubilizing Bacteria (PSB) for the character speed of germination which is regarded as very important indicator of seed vigour. Germination percentage for untreated control was 86.6% after one month of storage and found to be reduced to 76.67% after three month of storage but for treated *Azospirillum* in case of sonalika it was retained at 83.33%. V_1T_1 , V_2T_1 and V_3T_1 recorded improved germination even after six month of storage. The seedling characters like shoot and root length was greatly improved with T_1 and T_2 for almost the varieties included in this programme. The character vigour index greatly enhanced in case of V_1T_1 , V_1T_2 , V_3T_1 and V_3T_2 after one month and six months of storage. This work reported utilization of *Azospirillum* and PSB as seed inoculation for wheat varieties to enhance seed storability and seedling growth parameter. It can be concluded that, *Azospirillum* and PSB seed inoculation can be recommended for wheat for better seedling growth storability of seed

Keywords: *Azospirillum*, PSB, Seed inoculation, Varieties, Wheat

INTRODUCTION

Wheat (*Triticum aestivum* L.) can be classified as winter or spring growth habit based on flowering responses to cold temperatures. Biofertilizer like *Rhizobium*, *Azotobacter*, *Azospirillum* and blue green algae have been used for many years. *Azospirillum* inoculants are recommended mainly for wheat, sorghum, millets, maize, sugarcane and vegetable crops. Biofertilizer is defined as a substance which contains living organisms that when applied to seed, plant surface, or soil, colonize the rhizosphere or the interior of plant and promote growth by increasing supply or availability of primary nutrients to the host plant (Vessey, 2003). Biofertilizers are well recognized as an important component of integrated plant nutrient management for sustainable agriculture and hold a great promise to improve crop yield (Narula *et al.*, 2005). Plant growth promoting rhizobacteria are free living micro-organisms having beneficial effects on plants by colonizing their roots. They include such effects as the production of phytohormones; auxin, cytokinins and gibberellins (Garcia *et al.*, 2001), enhancing release of the nutrients (Nautiyal *et al.*, 2000). *Azotobacter* and *Azospirillum* have previously significantly increased

wheat and barley yield in irrigated as well as in rainfed crops (Pauw De *et al.*, 2008). To maintain the production potential as well as seed or grain quality of wheat, a concrete production technique is essential. One of the major essential elements for growth of plants is nitrogen. The Green Revolution (GR) technology adoption between 1960 to 2000 has increased wide varieties of agricultural crop yield per hectare which increased 12-13% food supply in developing countries. Nitrogen is required in large quantities for plants to grow, since it is the basic constituents of proteins and nucleic acids). Bio-fertilizers are the formulation of living organisms, which are able to fix atmospheric Nitrogen in the available from plants either by living frequency in soil or being associated symbiotically with plants (Subba Rao *et al.*, 1993). *Azospirillum*, a bacterial fertilizer is highly beneficial micro-organism for cereals, cotton, plantation crops and other crops. The main function of *Azospirillum* inoculation is to assimilate atmospheric nitrogen and fix in soil and finally providing the growing plant. The aim of present investigation was to study the effect of *Azospirillum* and PSB inoculation on different seed quality parameters like speed of germination, storability and different seedling parameters like fresh and dry weight, seedling length and vigour index.

MATERIALS AND METHODS

An experiment was carried out during the year of 2012-2013 to 2013-2014 in the department of Seed Science and Technology, Faculty of Agriculture, Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, Nadia, West Bengal. The experiment was done with three treatments and three replications with four varieties. The varieties were: V₁ (Sonalika), V₂ (PBW 443), V₃ (HD 2821) and V₄ (K 9107) and the treatments were: T₁ (*Azospirillum*), T₂ (PSB) and T₃ (Control) with thrice replicates. The data so obtained as described earlier were subjected to statistical analysis by Analysis of Variance method (Gomez and Gomez, 1984). The Standard Error mean (SEM±) and the value of critical difference (CD) were computed to compare the difference between means have been provided in the tables of results. The data on germination and field emergence were transformed into angular (arcsine) values (Snedecor and Cochran, 1967) transformed values are given in the parenthesis and then subject to statistical analysis.

The statistical calculations were prepared by Factorial Design for analysis of data linked to varietal consequence and Completely Randomized Design (CRD) for laboratory data. All statistical analyses were done using SPSS (version 10.0, 1990) on a desktop Computer. The initial seed moisture content was 12%. The seeds were stored in paper packet for one, three and six months of storage. The following seed quality parameters are recorded in the lab like Germination Percentage, Speed of germination, Root length, Shoot length, Seedling Fresh Weight, Dry Weight and Vigour index. The analysis of variance method (Cochran and Cox, 1963) was followed to analyze various data statistically. The significance of different sources of variation was tested by "Error Mean Square Method" of Fisher Snedecor's 'F' test at probability level 0.05. Standard error of Mean (SEM±) and critical difference (C.D.) have been estimated to compare the differences between means.

RESULTS AND DISCUSSION

The influence of different seed treatment on different varieties of wheat, particularly on the speed of germination which is regarded as very important indicator of seed vigour, found that the variety sonalika (V₁) was the best towards *Azospirillum* treatment (T₁) and in treatment in PSB (T₂), V₁ was best (Table 1). Speed of germination after one month of storage recorded highest volume in V₁T₁ and V₁T₂ than control. After three months of storage V₁T₁ and V₃T₂ responded better than control (Table 1). Response of different varieties after six months of storage was greatly pronounced in V₁T₁, V₁T₂, V₂T₂ and V₃T₂ for the character speed of germination (Table 1).

Germination percentage for control, after one month of storage was 86.67% which was found to be reduced to

76.67%, after three months of storage in case of V₁ but for T₁ it was retained at 83.33% after three months of storage (Table 2). V₁T₁, V₂T₁ and V₃T₁ recorded improved germination percentage than control. After six months of storage, control (T₃) exhibited loss of germination percentage to the extent of 63.33% in case of V₃ whereas the V₃T₁ and V₃T₂ maintained at 73.33% (Table 2). In case of soybean after 30 days plant maximum seed germination observed by biofertilizer treated plant (82.85%) and minimum by chemical fertilizer treated plants (0%) stated by Javed and Panwar (2013). Similar results were also reported by Strelec *et al.* (2010) to indicate significant influence of storage conditions on moisture content, germination and vigour changes during storage of wheat seeds, as well as varietal dependence of seed viability recorded. Response of different wheat varieties towards the character shoot length recorded high value for V₁T₁, V₁T₂ than control (Table 3). Similarly response was also obtained for V₂T₁, V₂T₂, V₃T₁ and V₃T₂ after three months of storage. After six months of storage, treatment effect towards shoot length was better for V₁T₁, V₁T₂, V₂T₁, V₂T₂, V₃T₁, V₃T₂ and V₄T₁ than control. Indicating *Azospirillum* and PSB seed treatment has the ability for improving seedling shoot length. Similar higher values were also recorded for V₁T₁, V₃T₁, V₃T₂ than control (Table 3). The biofertilizer resulted in maximum reduction of seed rot and foot/root rot (*Fusarium oxysporum*) of bushbean stated by Khalequzaman and Hossain (2008).

The character root length was greatly improved by T₁ and T₂ for V₁ but for V₃T₁ and V₄T₂ the results were lower than control. Enhanced root length was also found after six months of storage. Higher values were recorded for V₁T₁, V₁T₂ and V₄T₁ than control (Table 4). Chandrasekhar (2003) observed that the plant growth parameters viz., root length and number of leaves per plant in green gram plants at 45 Days were significantly increased due to inoculation of P-solubilizing fungal strains along with rock phosphate application as compared to rock phosphate alone (control).

Wheat varieties responded better towards seed treatment for the character fresh seedling weight after one month of storage for V₁T₁, V₁T₂, V₂T₁, V₂T₂, V₃T₁, V₃T₂ and V₄T₁, V₄T₂ than control (Table 5). After six months of storage, the seed treatment effect was observed for V₁T₁, V₁T₂, V₃T₁, V₃T₂ and V₄T₁, V₄T₂ but slightly reduced in V₂T₂. The variety-treatment interaction was greatly positive in V₁T₁, V₁T₂, V₃T₁, V₃T₂ and V₄T₁ and V₄T₂ (Table 5). Mahfouz and Sharafeldin (2007) have suggested the use of biofertilizers to increase the fresh weight of corn. The seedling dry weight, which is regarded as a great contributor towards seed vigour was also greatly enhanced by seed treatment. After one month of storage, V₁T₁, V₁T₂ recorded such type of enhancement response of variety was for to be best both after three and six months of

Table 1. Biofertilizers effect on speed of germination at after one, three and six months of storage (Three replicates).

V/T	Speed of germination											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	2.26	2.33	2.20	2.26	2.20	2.06	2.07	2.11	2.08	1.93	1.82	1.94
V2	1.78	2.27	2.30	2.12	1.63	2.14	2.29	2.02	1.49	2.08	1.94	1.84
V3	2.07	2.23	1.75	2.02	1.80	2.29	1.84	1.97	1.45	1.90	1.63	1.66
V4	2.07	2.30	2.45	2.27	2.00	2.09	2.29	2.13	1.66	2.04	2.06	1.92
Mean	2.04	2.28	2.18		1.91	2.14	2.12		1.67	1.99	1.86	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	0.072	0.063	0.125		0.028	0.024	0.049		0.023	0.020	0.039	
CD(P=0.05)	0.211	0.183	0.366		0.082	0.071	0.143		0.066	0.058	0.115	
CD(P=0.01)	0.836	0.724	1.447		0.326	0.283	0.565		0.263	0.228	0.455	

V₁-Sonalika, V₂- PBW 443, V₃, HD- 2821 and V₄- K 9107 and the treatments were- T₁-Azospirillum, T₂-PSB and T₃-Control.

Table 2. Biofertilizers effect on germination percentage at after one, three and six months of storage (Three replicates).

V/T	Germination %											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	96.67	96.67	86.67	93.33	83.33	83.33	76.67	81.11	76.67	73.33	70.00	73.33
V2	96.67	86.67	96.67	93.33	83.33	73.33	83.33	80.00	76.67	56.67	73.33	68.89
V3	96.67	93.33	93.33	94.44	83.33	83.33	76.67	81.11	73.33	73.33	63.33	70.00
V4	80.00	83.33	90.00	84.44	70.00	70.00	76.67	72.22	56.67	56.67	63.33	58.89
Mean	92.50	90.00	91.67		80.00	77.50	78.33		70.83	65.00	67.50	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	2.222	1.925	3.849		2.003	1.735	3.469		2.079	1.800	3.600	
CD(P=0.05)	6.486	5.617	11.234		5.847	5.063	10.127		6.067	5.254	10.509	
CD(P=0.01)	25.656	22.219	44.437		23.126	20.028	40.055		1.468	1.271	2.542	

V₁-Sonalika, V₂- PBW 443, V₃, HD- 2821 and V₄- K 9107 and the treatments were- T₁-Azospirillum, T₂-PSB and T₃-Control.

Table 3. Biofertilizers effect on shoot length (cm) at after one, three and six months of storage (Three replicates).

V/T	Shoot length (cm)											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	15.47	11.37	10.17	12.34	14.43	10.47	9.53	11.48	10.62	5.89	6.00	7.50
V2	15.67	14.99	12.27	14.31	12.93	12.50	11.73	12.39	8.72	8.20	7.22	8.05
V3	15.07	16.07	10.30	13.81	13.67	13.83	10.43	12.64	9.45	9.85	5.99	8.43
V4	14.00	9.34	9.09	10.81	12.30	8.90	9.00	10.07	8.19	5.37	5.33	6.30
Mean	15.05	12.94	10.46		13.33	11.43	10.18		9.25	7.33	6.14	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	0.282	0.245	0.489		0.125	0.109	0.217		0.127	0.110	0.220	
CD(P=0.05)	0.825	0.714	1.428		0.366	0.317	0.634		0.371	0.321	0.643	
CD(P=0.01)	3.261	2.824	5.649		1.447	1.253	2.506		1.468	1.271	2.542	

V₁-Sonalika, V₂- PBW 443, V₃, HD- 2821 and V₄- K 9107 and the treatments were- T₁-Azospirillum, T₂-PSB and T₃-Control.

Table 4. Biofertilizers effect on root length (cm) at after one, three and six months of storage (Three replicates).

V/T	Root length (cm)											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	13.36	13.43	11.38	12.72	11.93	12.53	10.83	11.77	10.87	11.47	9.00	10.44
V2	15.03	13.83	14.87	14.58	12.80	12.80	13.33	12.98	11.67	11.80	12.03	11.83
V3	12.42	13.47	13.00	12.96	12.20	12.93	12.27	12.47	10.30	10.33	10.67	10.43
V4	15.40	10.17	11.09	12.22	13.80	9.93	9.97	11.23	12.60	8.30	8.33	9.74
Mean	14.05	12.73	12.58		12.68	12.05	11.60		11.36	10.48	10.01	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	0.207	0.179	0.358		0.165	0.143	0.286		0.124	0.107	0.214	
CD(P=0.05)	0.604	0.523	1.045		0.482	0.418	0.835		0.361	0.313	0.626	
CD(P=0.01)	2.387	2.068	4.135		1.907	1.652	3.303		1.428	1.237	2.474	

V₁-Sonalika, V₂- PBW 443, V₃, HD- 2821 and V₄- K 9107 and the treatments were- T₁-Azospirillum, T₂-PSB and T₃-Control.

storage (Table 6). The similar work was done in case of corn and showed the biofertilizers increased the dry weight proposed by Mahfouz and Sharaf eldin (2007). The study revealed that maximum germination (92%), viability (95%), germination speed (1.58), germination

energy (70.0%) were recorded in polybags under complete darkness in *Aconitum heterophyllum* wall ex. Royle while minimum were recorded under partial light in cloth bags. Similarly, in case of *Podophyllum hexandrum* same results were obtained stated by

Table 5. Biofertilizers effect on fresh weight (g) at after one, three and six months of storage (Three replicates).

V/T	Fresh weight (g)											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	1.21	1.30	0.86	1.12	1.07	1.17	0.79	1.01	1.06	1.14	0.83	1.01
V2	0.74	0.73	0.71	0.73	0.69	0.66	0.67	0.67	0.67	0.65	0.67	0.66
V3	1.15	1.13	0.88	1.05	1.17	1.17	0.82	1.05	1.10	1.07	0.85	1.01
V4	0.93	0.90	0.80	0.88	0.92	0.85	0.76	0.85	0.85	0.83	0.74	0.81
Mean	1.01	1.01	0.81		0.96	0.96	0.76		0.92	0.92	0.77	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	0.027	0.024	0.047		0.015	0.013	0.027		0.021	0.018	0.036	
CD(P=0.05)	0.080	0.069	0.138		0.045	0.039	0.078		0.060	0.052	0.104	
CD(P=0.01)	0.316	0.273	0.547		0.178	0.154	0.309		0.238	0.206	0.411	

V₁-Sonalika, V₂- PBW 443, V₃-HD- 2821 and V₄- K 9107 and the treatments were- T₁-*Azospirillum*, T₂- PSB and T₃- Control.

Table 6. Biofertilizers effect on dry weight (g) at after one, three and six months of storage (Three replicates).

V/T	Dry weight (g)											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	0.18	0.19	0.13	0.17	0.17	0.17	0.10	0.15	0.14	0.15	0.07	0.12
V2	0.12	0.13	0.41	0.22	0.11	0.10	0.10	0.10	0.06	0.06	0.04	0.05
V3	0.17	0.13	0.13	0.14	0.17	0.12	0.12	0.14	0.13	0.09	0.07	0.09
V4	0.13	0.11	0.13	0.12	0.12	0.10	0.13	0.12	0.06	0.05	0.07	0.06
Mean	0.15	0.14	0.20		0.14	0.12	0.11		0.09	0.08	0.06	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	0.049	0.042	0.084		0.002	0.002	0.003		0.009	0.008	0.016	
CD(P=0.05)	0.142	0.123	0.247		0.005	0.004	0.009		0.027	0.024	0.047	
CD(P=0.01)	0.563	0.488	0.975		0.020	0.017	0.035		0.108	0.094	0.187	

V₁-Sonalika, V₂- PBW 443, V₃-HD- 2821 and V₄- K 9107 and the treatments were- T₁-*Azospirillum*, T₂- PSB and T₃- Control.

Table 7. Biofertilizers effect on Vigour index at after one, three and six months of storage.

V/T	Vigour index											
	After one month of storage				After three months of storage				After six months of storage			
	T1	T2	T3	Mean	T1	T2	T3	Mean	T1	T2	T3	Mean
V1	2785	2402	1868	2352	2198	1915	1562	1892	1648	1270	1049	1322
V2	2965	2498	2624	2696	2146	1856	2088	2030	1563	1132	1409	1368
V3	2660	2757	2172	2530	2154	2230	1739	2041	1449	1482	1053	1328
V4	2347	1631	1812	1930	1821	1318	1454	1531	1179	774	865	939
Mean	2689	2322	2119		2080	1830	1711		1459	1165	1094	
	V	T	VXT		V	T	VXT		V	T	VXT	
S.Em(±)	64.507	55.865	111.730		46.73	40.47	80.94		36.57	31.67	63.35	
CD (P=0.05)	188.28	163.06	326.12		136.4	118.1	236.2		106.8	92.45	184.9	
CD (P=0.01)	744.75	644.97	1289.94		539.5	467.2	934.4		422.3	365.7	731.4	

V₁-Sonalika, V₂- PBW 443, V₃-HD- 2821 and V₄- K 9107 and the treatments were- T₁-*Azospirillum*, T₂- PSB and T₃- Control.

Verma *et al.* (2011). A storage experiment was conducted to understand the effect of organic and integrated management practices of seed production and storage containers along with organic (insecticide and fungicide) and organic (botanicals) as seed treatments on seed viability of scented rice cv. Mugad sugandha for 20 months reported by Raikar *et al.* (2011). Chattha *et al.* (2012) observed the effect of different packing materials (metal bin, earthen bin, plastic bag, cloth bag and gunny bag) and grain moisture content at packing (10% and 16%) on viability of wheat seed for ten months of storage.

The seedling quality parameter, vigour index, where two important seed quality parameters like germination percentage and seedling length was taken into account to find out the response of varieties under study towards maintaining or enhancing vigour status after six months of storage. It is noted that, vigour index was greatly enhanced in case of V₁T₂, V₁T₃, V₃T₁, V₃T₂ after one, three and six months of storage. After six

months of storage, in comparison to control, all the treatments recorded maximum value except V₄T₂ and V₂T₂. Other V₁T₁, V₁T₂, V₂T₁, V₃T₁, V₃T₂ and V₄T₁ all recorded better value than control. Similar results were reported by Albrecht *et al.* (1981), Mishra *et al.* (1998) and Rout *et al.* (2001) in case of maize. The highest germination, normal seedlings and vigour index which were followed by polythene bag, where Gunny bag showed the lowest germination, normal seedlings and vigour index upto 60 days after storage by Khalequzaman *et al.* (2012).

Conclusion

From findings it was revealed that, *Azospirillum* and PSB seed inoculation can be recommended for wheat for better seedling growth storability of seed. It should also be noted that, these bio-fertilizers are less expensive than inorganic manures and they are also eco-friendly. More number of wheat genotypes may be used in future for recommendation of actual bio-

fertilizers dose for yield enhancement.

REFERENCES

- Albrecht, S.L., Okon, Lonquist L and Burries, R.H. (1981). Nitrogen fixation by corn *Azospirillum* associations in a temperature climate. *Crop Science*, 21: 301-306.
- Chandrasekhar, B.S. (2003). Studies on mineral phosphate solubilizing fungi from vertisols of Northern Karnataka and other Biofertilizer potential. Ph.D. Thesis, Univ. Agric. Sci., Dharwad, Karnataka, India.
- Chattha, S.H., Jamali, L.A., Ibutoto, K.A. and Mangio, H.R. (2012). Effect of different packing materials and Storage conditions on the viability of wheat Seed (TD-1 variety). *Sci. Tech. and Dev.*, 31 (1): 10-18.
- Cochran, W.G and Cox, G.M. (1963). Experimental Designs. In: 1st India Ed. Asia publishing House, Bombay, Calcutta, New Delhi and Madras.
- Garcia, de Salamone, I.E. Hynes, R.K. and Nelson, L. M. (2001). Cytokinins production by plant growth promoting rhizobacteria and selected mutants. *Can. J. Microbiol.*, 47: 404-411.
- Gomez, K.A. and Gomez, A.A. (1984). Statistical Procedures for Agricultural Research, 2nd edn. Singapore: John Wiley & Sons.
- Javed, S. and Panwar, A. (2013). Effect of biofertilizer, vermicompost and chemical fertilizer on different biochemical parameters of Glycine max and Vigna mungo. *Recent Research in Science and Technology*. 5 (1): 40-44.
- K.M. Khalequzaman and I. Hossain (2008). Effect of Seed Treatment with Rhizobium Strains and Biofertilizers on foot/root rot and Yield of Bushbean in *Fusarium oxysporum* infested soil. *J. Agric. Res.*, 46 (1): 55-65
- Khalequzaman, K.M., Rashid, M.M., Hasan, M.A. and Reza, M. M. A. (2012). Effect of storage containers and storage periods on the seed quality of french bean (*Phaseolus vulgaris*). *Bangladesh J. Agril. Res.*, 37 (2): 195-205.
- Mahfouz S.A. and Sharaf-Eldin M.A. (2007). Effect of mineral vs. biofertilizer on growth, yield and essential oil content of fennel (*Foeniculum vulgare* Mill.). *Int Agro-phys.* 21: 361 -366.
- Mishra, M.; Patyonshi, A.K. and Jena, D. (1998). Effect of biofertilizers on productivity of maize (zea mays), *Indian Journal of Agronomy*. 43: 307-310.
- Narula, N. Kumar, V. Singh, B. Bhatia, R. and Lakshminarayana, K. (2005). Impact of Biofertilizer on rain yield in spring wheat under varying fertility condition and wheat cotton rotation. *Archiv. Agron. and Soil Sci.*, 51: 79-89.
- Nautiyal, C.S. Bhadauria, S. Kumar, P.L. Momdal, R. and Verma, D. (2000). Stress induced phosphate solubilization in bacteria isolated from alkaline soils. *FEMS Micro-biology Lett.* 182-:291-296.
- Pauw, De. Mirghasemi, E.A. Ghaffari, A. and Nseir, B. (2008). Agro ecological zones of Karkheh River, Basin. Research Report, ICARDA, Aleppo, Syria.
- Raika, S.D., Vyakarnahal, B.S., Biradar, D. P., Deshpande, V.K. and Janagoudar, B.S. (2011). Effect of seed source, containers and seed treatment with chemical and biopesticide on storability of scented rice Cv. Mugad sugandha. *Karnataka J. Agric. Sci.*, 24 (4): 448-454.
- Rout, D., Satalathy, M.R.; and Mohapatra, B.K. (2001). Effect of biofertilizers on nitrogen economy in maize. *Madras Agricultural Journal*. 88:7-9.
- Snedecor, G and Cochran, W.G. 1967. Statistical methods, Oxford and IBH Publishing Company, Bombay, pp. 135 -197.
- Strelec, I., Popovic, R., Ivanisic, I., Jurkovic, V., Jurkovic, Z., Hardi, Z.U. and Sabo, M. (2010). Influence of temperature and relative humidity on grain moisture, germination and vigour of three wheat cultivars during one year storage. *Poljoprivreda*, 16 (2): 20-24.
- Subba Rao, N.S. (1993). Biofertilizers in Agriculture and Forestry, Oxford and IBM publishing co. (p) Ltd. 3 rd edition.
- Verma, N.K., Verma, A.K. and Kumar, D. (2011). Effect of storage containers on seed germination and viability in *Aconitum heterophyllum* and *Podophyllum hexandrum* – endangered medicinal plants species of Himalayan region. *Biological Forum – An International Journal*, 1 (1): 8-11.
- Vessey, J.K. (2003). Plant growth promoting rhizobacteria as biofertilizer. *Plant and Soil*. 255: 571-586.